

UNIVERSAL EQUIVALENCE OF PARTIALLY COMMUTATIVE METABELIAN LIE ALGEBRAS

E. N. POROSHENKO, E. I. TIMOSHENKO

ABSTRACT. In this paper, we find a criterium for universal equivalence of partially commutative Lie algebras whose defining graphs are trees. Besides, we obtain bases for partially commutative metabelian Lie algebras.

1. INTRODUCTION

Let $G = \langle X, E \rangle$ be an undirected graph without loops with the finite set of vertices $X = \{x_1, \dots, x_n\}$ and the set of edges E ($E \subseteq X \times X$). We denote the elements of E by $\{x, y\}$.

Consider a variety $\mathfrak{M}$ of Lie algebras over a ring R . A *partially commutative Lie algebra in $\mathfrak{M}$ with a defining graph G* is a Lie algebra $\mathcal{L}_R(X; G)$ defined as

$$\mathcal{L}_R(X; G) = \langle X \mid [x_i, x_j] = 0 \iff \{x_i, x_j\} \in E; \mathfrak{M} \rangle$$

in $\mathfrak{M}$. Thus, in this algebra, the variety identities and the defining relations hold together. If there is no ambiguity denote this algebra just by $\mathcal{L}(X; G)$. For simplicity (to avoid using the notation for the set of edges), we write $\{x_i, x_j\} \in G$ instead of $\{x_i, x_j\} \in E$.

Usually, the varieties whose identities do not imply additional relations of vertices' commutation are considered. It means that two vertices commute if and only if they are adjacent in G . In this paper, we study partially commutative metabelian Lie algebras. These algebras clearly possess the property indicated above. It follows, for example, from the structure of the bases for partially commutative metabelian Lie algebras (see Theorem 3.3).

Along with the variety of Lie algebras, one can consider other varieties of algebras and groups. The most actively studied objects of this kind are partially commutative groups which are defined by commutativity relations in the variety of all groups. Some papers (see, [4, 7, 5] for example), are devoted to universal theories of partially commutative metabelian groups. In [2] and [6], partially commutative associative algebras were studied.

This paper is organized as follows. In Sec. 2, preliminary definitions and results are given.

In Sec. 3, we find bases for partially commutative metabelian Lie (see Theorem 3.3). This theorem is used a great deal in the paper but it is also interesting in itself.

In Sec. 4, we prove Theorem 4.7 and Theorem 4.9 which give information on the centralizers of some elements and on the annihilators of some elements in the derived subalgebra of partially commutative metabelian Lie algebras. We need these results for the study of the universal theories of partial commutative metabelian Lie algebras.

The main result of the paper is Theorem 5.9 which is proved in Sec. 5. This is a criterium for coincidence of the universal theories of partially commutative metabelian Lie algebras whose defining graphs are trees. It is easy to verify if this

condition holds. So, the problem of universal equivalence for the algebras defined above is algorithmically solvable.

2. PRELIMINARIES

Let $M(X)$ denote the free metabelian Lie R -algebra with the set of generators X . A *partially commutative metabelian Lie algebra* generated by X with the defining graph G is the Lie algebra $M(X; G) = M(X)/I$, i.e. this is the Lie algebra, defined by the set of relations

$$(1) \quad [x_i, x_j] = 0, \text{ if } \{x_i, x_j\} \in G$$

in the variety of metabelian Lie algebras.

Definition 2.1. Let G be a graph. A vertex of G is called an *endpoint* if its degree is equal to 1.

In the paper, we need a couple results with reference to free metabelian Lie algebras. Let us formulate these results here.

Theorem 2.2. [8] *Let L be the free polynilpotent Lie ring corresponding to the sequence $n_1, \dots, n_k$, ($k \geq 2$). If $x, y \in L$ are such that $[x, y] = 0$ then either x and y are linearly dependent or $x, y \in L_{n_1, \dots, n_k}$.*

We need this statement only in the case of a free metabelian Lie algebra. It is easy to see that Theorem 2.2 also holds for a free metabelian Lie algebras over any integral domain.

Usually, to study an algebra, it is very useful to know its basis. In [1, 8], linear bases for free polynilpotent Lie algebras were found. Since we need only the structure of a basis for metabelian Lie algebras, we give the description of a basis in this case only.

Theorem 2.3. *A basis of the free metabelian Lie algebra $M(X)$ consists of the set X together with Lie monomials of the form*

$$(2) \quad [\dots [x_{i_1}, x_{i_2}], \dots, x_{i_m}], \quad \text{where } m \geq 2, \quad x_{i_2} < x_{i_1}, \quad x_{i_2} \leq x_{i_3} \leq \dots \leq x_{i_m}.$$

Let us denote this basis by $\mathfrak{B}(X)$. Since it consists of left-normed Lie monomials only, in this paper, we omit all Lie brackets except the outer pair, i.e. we write $[x_{i_1}, x_{i_2}, \dots, x_{i_m}]$ instead of $[\dots [x_{i_1}, x_{i_2}], \dots, x_{i_m}]$.

By definition, we suppose $\ell([x_{i_1}, x_{i_2}, \dots, x_{i_r}]) = r$ for any left-normed Lie monomial and say that r is the *length* of the monomial $[x_{i_1}, x_{i_2}, \dots, x_{i_r}]$.

Definition 2.4. Let u be a Lie monomial. The *multidegree* of u is the vector $\bar{\delta} = (\delta_1, \delta_2, \dots, \delta_n)$, where δ_i is the number of occurrences of x_i in u .

Let us denote by $\text{mdeg}(u)$ the multidegree of u . If all monomials of a Lie polynomial g have the same multidegree $\bar{\delta}$ then we call such polynomial *homogeneous* and write $\text{mdeg}(g) = \bar{\delta}$.

Note that we can define the sum of multidegrees as the sum of the corresponding vectors. We also use the notation $\text{mdeg}_i(u)$ for the number of x_i in u , i.e. for the i th coordinate of $\text{mdeg}(u)$.

Let us define an order on $\mathbb{Z}^n$, namely for arbitrary vectors $\bar{\delta} = (\delta_1, \dots, \delta_n)$ and $\bar{\gamma} = (\gamma_1, \dots, \gamma_n)$, let us set $\bar{\delta} > \bar{\gamma}$ if there is k such that $\delta_k > \gamma_k$ for some k but $\delta_i = \gamma_i$ for $i > k$. We use this order to compare multidegrees. Define an order on $\mathfrak{B}(X)$. Let $u, v \in \mathfrak{B}(X)$. We say that u is *greater* than v , if one of the following conditions holds:

- (1) $\text{mdeg}(u) > \text{mdeg}(v)$;
- (2) $\text{mdeg}(u) = \text{mdeg}(v)$ and u is greater than v lexicographically.

We call this order by *the standard order* or just *the order* and denote it by " $\geq$ ". Let f be an arbitrary Lie polynomial. Denote by $\bar{f}$ the largest monomial (in the sense of the standard order) appearing in f without a coefficient by it. So, $f = \alpha\bar{f} + \sum_i \alpha_i u_i$, where $\alpha, \alpha_i \neq 0$ and $\bar{f} > u_i$ for all u_i .

Let $v = [x_{i_1}, \dots, x_{i_k}]$. It is easy to see that any permutation of the letters $x_{i_3}, \dots, x_{i_k}$ gives us a monomial equal to v in $M(X)$. Indeed,

$$\begin{aligned} & [x_{i_1}, x_{i_2}, \dots, x_{i_{s-1}}, x_{i_s}, x_{i_{s+1}}] \\ &= [[x_{i_1}, x_{i_2}, \dots, x_{i_{s-1}}], [x_{i_s}, x_{i_{s+1}}]] + [x_{i_1}, x_{i_2}, \dots, x_{i_{s-1}}, x_{i_{s+1}}, x_{i_s}] \\ &= 0 + [x_{i_1}, x_{i_2}, \dots, x_{i_{s-1}}, x_{i_{s+1}}, x_{i_s}]. \end{aligned}$$

Consequently,

$$[x_{i_1}, x_{i_2}, \dots, x_{i_{s-1}}, x_{i_s}, x_{i_{s+1}}, \dots, x_{i_k}] = [x_{i_1}, x_{i_2}, \dots, x_{i_{s-1}}, x_{i_{s+1}}, x_{i_s}, \dots, x_{i_k}].$$

So, interchanging letters x_{i_s} and $x_{i_{s+1}}$ ($s \geq 3$) in v gives us an element equal to v in $M(X)$. We are left to note that the set of all transpositions permuting two neighbor elements generates the symmetric group on the set of these elements. Therefore, we can obtain any permutation of the letters $x_{i_3}, \dots, x_{i_k}$ in v interchanging pairs of neighbor elements.

$$(3) \quad [x_{i_1}, x_{i_2}, x_{i_3}, \dots, x_{i_k}] = [x_{i_1}, x_{i_2}, x_{\sigma(i_3)}, \dots, x_{\sigma(i_k)}],$$

where σ is a permutation on $\{i_3, i_4, \dots, i_k\}$.

Let $R[X]$ be the set of all commutative associative polynomials over R . It follows from the last paragraph that the derived subalgebra $M'(X)$ of $M(X)$ is an $R[X]$ -module with respect to the adjoint representation. Denote by $u.f$ the element of $M'(X)$ obtained by acting the element $f \in R[X]$ on $u \in M'(X)$. Namely, let us define $u.f$ inductively:

- (1) $u.y = [u, y]$ for any $y \in X$;
- (2) Let $f = y_1 y_2 \dots y_m$ for $m \geq 2$ and let $f_0 = y_1 y_2 \dots y_{m-1}$ then $u.f = (u.f_0).y_m$;
- (3) Finally, if $f = g + s$, where s is a commutative associative monomial then $u.f = u.g + u.s$.

For any monomial in $R[X]$ we also can define its multidegree as follows: $\text{mdeg}(\alpha x_1^{\gamma_1} x_2^{\gamma_2} \dots x_n^{\gamma_n}) = (\gamma_1, \gamma_2, \dots, \gamma_n)$, if $\alpha \neq 0$.

Let us remind the following well-known result. Let g be a non-associative polynomial in X and let $g = \sum_{\bar{\delta}} g_{\bar{\delta}}$, where $g_{\bar{\delta}}$ is a homogeneous polynomial such that $\text{mdeg}(g_{\bar{\delta}}) = \bar{\delta}$. If $g = 0$ in $M(X; G)$ then $g_{\bar{\delta}} = 0$ in this algebra for all $\bar{\delta}$. It follows from homogeneity of Lie algebras, metabelian algebra identities, and partial commutativity relations. Indeed, if $g = 0$ then this polynomial can be nullified by using a finite set of identities and relations indicated above. Each step moves a homogeneous polynomial to homogeneous one of the same multidegree. Therefore, summands of different homogeneous components cannot cancel with each other.

Then, we introduce some homomorphisms on $M(X; G)$. First, let us remind a well-known fact. Let $X = \{x_1, \dots, x_n\}$ be the set of generators of a Lie algebra $\mathcal{L}_1$. Consider a map $\varphi : X \rightarrow \mathcal{L}_2$, where $\mathcal{L}_2$ is also a Lie algebra. If all identities and relations of $\mathcal{L}_1$ hold under φ then this map can be extended to a homomorphism from $\mathcal{L}_1$ to $\mathcal{L}_2$ uniquely. So, all maps defined below are homomorphisms.

Let X be the set of generators of the partially commutative metabelian Lie algebra $M(X; G)$. For any non-empty subset Y of X , denote by G_Y the subgraph of G generated by Y .

Definition 2.5. The *projection* from $M(X; G)$ onto $M(Y; G_Y)$ is a homomorphism $\pi_Y : M(X; G) \rightarrow M(Y; G_Y)$ defined on X as follows:

$$(4) \quad \pi_Y(x_i) = \begin{cases} x_i, & \text{if } x_i \in Y \\ 0, & \text{if } x_i \notin Y. \end{cases}$$

Definition 2.6. Let $\overline{G}$ be obtained from G by adding some edges. An *identical simplification* from $M(X; G)$ onto $M(X; \overline{G})$ is the homomorphism $\varphi : M(X; G) \rightarrow M(X; \overline{G})$ defined on the set of generators identically (i.e. $\varphi(x_i) = x_i$ for $x_i \in X$).

Let G be an arbitrary graph, $G_1, G_2, \dots, G_r$ its connected components, and X_i the set of the vertices of G_i for $i = 1, 2, \dots, r$. Given t such that $0 \leq t < r$, define the sets $X' = \bigcup_{i=1}^t X_i$, $X'' = \bigcup_{j=t+1}^r X_j$. Consider a graph $\tilde{G}$ obtained from the subgraph $G_{X'}$ adding some set Z of isolated vertices (Z may be empty). By Y denote the set of vertices of $\tilde{G}$. In other words, $Y = X' \cup Z$.

Definition 2.7. An *identification on connected components* $G_{t+1}, \dots, G_r$ is a homomorphism $\varphi : M(X; G) \rightarrow M(Y; \tilde{G})$, acting on X' identically and mapping all vertices in $X_k \subseteq X''$ ($k = t+1, \dots, r$) to multiples of a fixed element $y_k \in Y$ (i.e. to elements of the form αy_k , where $\alpha \in R$ can be different for different vertices in X_k). If $X' = \emptyset$ (this means $t = 0$) then the identification of connected components is called *complete*.

Definition 2.7 implies that if an identification of connected components is complete then $\tilde{G}$ is totally disconnected, therefore $M(Y; \tilde{G}) = M(Y)$.

Finally, let us recall some terminology related to universal theories of algebraic systems.

Definition 2.8. An $\exists$ -sentence is a formula without free variables such that it is of the form

$$\exists w_1 \dots w_m \Phi(w_1, \dots, w_m),$$

where $\Phi(w_1, \dots, w_m)$ is a formula of predicate calculus in the corresponding algebraic system such that this formula does not contain quantifiers.

Definition 2.9. The set of all $\exists$ -sentences that are true in a Lie algebra L is called the *existential theory* or the $\exists$ -theory of this Lie algebra.

Definition 2.10. Lie algebras are called *existentially equivalent* if their existential theories coincide.

The notion of *universal theory* (or $\forall$ -theory) of a Lie algebra is defined analogously as well as the notion of universal equivalence of Lie algebras.

It is easy to see that Lie algebras L_1 and L_2 are existentially equivalent if and only if these Lie algebras are universally equivalent.

The procedure of exchanging functional symbols by predicate ones is well-known in model theory. Any set with all predicates induced on it is a submodel.

Let us formulate a well-known result in model theory.

Theorem 2.11. *Arbitrary algebraic systems (ex., Lie algebras) L_1 and L_2 are universally equivalent if and only if each finite model of the first algebraic system is isomorphic to a finite model of the second one.*

3. BASES OF PARTIALLY COMMUTATIVE METABELIAN ALGEBRAS

Given $u = [x_{i_1}, x_{i_2}, \dots, x_{i_k}]$, let X_u be the set of all generators appearing in u . Let's denote by G_u the subgraph of G , generated by the set X_u . Similarly, we define X_g and G_g for a homogeneous Lie polynomial g , i.e. suppose that $X_g = X_u$ and $G_g = G_u$, where u is a monomial of g .

Lemma 3.1. *Suppose $u = [x_{i_1}, x_{i_2}, \dots, x_{i_k}]$, x_{i_1} and x_{i_j} are vertices belonging to the same connected component of G_u , and u' is obtained from u by interchanging x_{i_1} and x_{i_j} . Then $u = u'$ in $M(X; G)$.*

Proof. Since x_{i_1} and x_{i_j} are in the same connected component of G_u , there is a simple path $(x_{i_1}, y_1, y_2, \dots, y_s, x_{i_j})$ in this graph, i.e. a path consisting of elements of X_u all vertices of which are different. Let us proceed by induction on the number of intermediate vertices on such path (i.e. by induction on s).

If $s = 0$ then x_{i_1} and x_{i_j} are adjacent. Consider two cases. For $j = 2$ we have $[x_{i_1}, x_{i_2}, \dots, x_{i_k}] = [0, x_{i_3}, \dots, x_{i_k}] = 0$ and, analogously, $[x_{i_2}, x_{i_1}, \dots, x_{i_k}] = [0, x_{i_3}, \dots, x_{i_k}] = 0$.

Let $j \geq 3$. By (3), without loss of generality it can be assumed that $j = 3$. Since $[x_{i_1}, x_{i_3}] = 0$ we obtain

$$\begin{aligned} [x_{i_1}, x_{i_2}, x_{i_3}, x_{i_4}, \dots, x_{i_k}] &= [x_{i_1}, x_{i_3}, x_{i_2}, x_{i_4}, \dots, x_{i_k}] + [\dots [x_{i_1}, [x_{i_2}, x_{i_3}]], \dots, x_{i_k}] \\ &= -[x_{i_2}, x_{i_3}, x_{i_1}, x_{i_4}, \dots, x_{i_k}] \\ &= [x_{i_3}, x_{i_2}, x_{i_1}, x_{i_4}, \dots, x_{i_k}]. \end{aligned}$$

Next, let us suppose that the statement holds for all $l < s$. Consider the path $(x_{i_1}, y_1, \dots, y_s, x_{i_j})$ from x_{i_1} to x_{i_j} in G_u . As it was shown above, the monomial u_1 obtained by interchanging x_{i_1} and y_1 in u is equal to u in $M(X; G)$. Therefore,

$$[x_{i_1}, x_{i_2}, \dots, y_1, \dots, x_{i_k}] = [y_1, x_{i_2}, \dots, x_{i_1}, \dots, x_{i_k}]$$

in $M(X; G)$. We have $(y_1, \dots, y_s, x_{i_j})$ contains $s - 1$ intermediate vertices. So, by inductive assumption, $u_2 = u_1$, in $M(X; G)$ where u_2 is the monomial obtained by interchanging y_1 and x_{i_j} in u_1 .

Finally, to obtain the monomial u' we are left to interchange x_{i_1} and y_1 in u_2 . If either y_1 or x_{i_1} is the second letter in u_2 then one can interchange the first two letters by the anticommutativity identity, after that interchange y_1 and x_{i_1} by the induction hypothesis, and again interchange the first two letters. If neither y_1 nor x_{i_1} is the second letter of u_2 then $u_2 = u'$ in $M(X; G)$ by (3). Therefore, $u = u'$ in $M(X; G)$. $\square$

Lemma 3.2. *Let $u = [x_{i_1}, \dots, x_{i_k}]$ be a monomial of the form (2). Then u is equal to zero in $M(X; G)$ if and only if the vertices x_{i_1} and x_{i_2} are in the same connected component of G_u .*

Proof. If x_{i_1} and x_{i_2} are in the same connected component of G_u , then the statement follows from Lemma 3.1.

Conversely, let x_{i_1} and x_{i_2} be in different connected components of G_u . We need to show that $u \neq 0$ in this case.

Let π_{X_u} be the projection of the algebra $M(X; G)$ onto the algebra $M(X_u, G_u)$. Since π_{X_u} acts identically on the set X_u we have $\pi_{X_u}(u) = u$.

Consider a set Y such that it contains by one element from each connected component of G_u and both x_{i_1} and x_{i_2} are in Y .

Define the complete identification of connected components $\psi : M(X_u; G_u) \rightarrow M(Y)$ mapping each element in X_u to the element in Y from the same connected component of G_u .

Applying ψ to the monomial u in $M(X_u; G_u)$, we obtain

$$0 = \psi([x_{i_1}, x_{i_2}, \dots, x_{i_k}]) = [x_{i_1}, x_{i_2}, \psi(x_{i_3}), \dots, \psi(x_{i_k})].$$

Since x_{i_2} is the smallest letter of u it is also the smallest letter in $\psi(u) = [x_{i_1}, x_{i_2}, \psi(x_{i_3}), \dots, \psi(x_{i_k})]$. By (3), one can permute the letters $\psi(x_{i_3}), \dots, \psi(x_{i_k})$ in $\psi(u)$ in such a way that they are in the non-descending order in the obtained monomial. The obtained monomial is of the form (2) and it is equal to $\psi(u)$ in

$M(Y)$. Therefore, it cannot be equal to 0 in this algebra. Consequently, $u \neq 0$ in $M(X; G)$ either. $\square$

Let us consider the algebra $M(X; G)$. By $B_{\bar{\delta}}^L(X)$ denote the set of all monomials of the form (2) whose multidegrees are equal to $\bar{\delta}$. Let us eliminate from $B_{\bar{\delta}}^L(X)$ all monomials u such that $u = 0$ in $M(X; G)$ (by Lemma 3.2, we should exclude all monomials $u = [x_{i_1}, x_{i_2}, \dots, x_{i_k}]$ such that x_{i_1} and x_{i_2} are in the same connected component of the graph G_u). By $B_{\bar{\delta}}''(X; G)$ denote the obtained set.

Given monomials $u_1 = [x_{i_1}, x_{i_2}, \dots, x_{i_k}]$ and $u_2 = [x_{j_1}, x_{j_2}, \dots, x_{j_k}]$ in $B_{\bar{\delta}}''(X; G)$ we write $u_1 \sim u_2$ if x_{i_1} and x_{j_1} are in the same connected component of $G_{u_1} = G_{u_2}$. By Lemma 3.1, it follows that if $u_1 \sim u_2$ then $u_1 = u_2$ in $M(X; G)$. Besides, it follows from (2) that $x_{i_2} = x_{j_2}$. The relation “ $\sim$ ” is obviously an equivalence relation. Consequently, there is the decomposition of $B_{\bar{\delta}}''(X; G)$ by the equivalence classes. Let us choose by one element from each equivalent class in such a way that the first letter of each chosen element is the the largest one among the first letters of all monomials in this equivalence class. Denote this set by $B_{\bar{\delta}}(X; G)$ (in particular, this set can be empty for some multidegrees).

Let us set

$$\mathfrak{B}(X; G) = \bigcup_{\bar{\delta}} B_{\bar{\delta}}(X; G),$$

where the union is taken by all multidegrees.

The following theorem holds.

Theorem 3.3. *The set $\mathfrak{B}(X; G)$ is a basis of the partially commutative metabelian Lie algebra $M(X; G)$.*

Proof. Since all elements of the same equivalence class are equal to each other in $M(X; G)$ and since $\mathfrak{B}(X; G)$ contains elements from all equivalence classes of the set $\bigcup_{\bar{\delta}} B_{\bar{\delta}}''(X; G)$ the algebra $M(X; G)$ is spanned by $\mathfrak{B}(X; G)$.

We are left to show that the elements of $\mathfrak{B}(X; G)$ are linearly independent. Let

$$(5) \quad g = \sum_j \alpha_j u_j,$$

where $u_j = [x_{j_1}, x_{j_2}, \dots, x_{j_{k_j}}] \in \mathfrak{B}(X; G)$, $\alpha_j \in R$. Suppose that $g = 0$ in $M(X; G)$. Without loss of generality (see Sec. 2) we may assume that g is a homogeneous polynomial. In particular, all k_j are equal to each other. For this reason, we write k instead of k_j . Note that the second letter in all monomials u_j is same, namely, it is the smallest letter appearing in u_j . Let us denote this letter by x .

By construction of $B_{\bar{\delta}}(X; G)$, the first letters of different monomials u_j in decomposition (5) belong to different connected components of G_g . Consider the set of the first letters of the monomials u_j appearing in (5). Extend this set to the set Y such that Y contains by one letter from each connected component of G_g and $x \in Y$. It can be done because the first letter of a non-zero monomial u_j cannot belong to the same connected component as x . So, x is the least element of Y .

Consider the projection $\pi_{X_g} : M(X; G) \rightarrow M(X_g; G_g)$. Since π_{X_g} acts identically on X_g we have $\pi_{X_g}(g) = g$.

Let $\psi : M(X_g; G_g) \rightarrow M(Y)$ be the complete identification of connected components mapping each element of X_g to the element of Y belonging to the same connected component of G_g .

Applying the map ψ to (5) we get

$$\begin{aligned} 0 &= \sum_j \alpha_j \psi(u_j) \\ &= \sum_j \alpha_j [\psi(x_{j_1}), \psi(x), \psi(x_{j_3}), \dots, \psi(x_{j_k})]. \end{aligned}$$

It was shown above that $\psi(x_{j_s}) \in Y$. If it is necessary we can permute the letters $\psi(x_{i_3}), \dots, \psi(x_{i_k})$ in each $\psi(u_j)$ in such a way that they are in the non-descending order in the obtained monomial. We obtain

$$\sum_j \alpha_j [x_{j_1}, x, y_{j_3}, \dots, y_{j_k}] = 0$$

in the free metabelian algebra $M(Y)$. Since x_{j_1} and x are in different connected components, $\psi(u_j) \neq 0$ in $M(Y)$. By Theorem 2.3, the set of monomials $\psi(u_j)$ is linearly independent in $M(Y)$. Consequently, $\alpha_j = 0$ for all j . We have proved that the monomials u_j are linearly independent in $M(X_g; G_g)$ so are they in $M(X; G)$. This completes the proof. $\square$

4. ANNIHILATORS AND CENTRALIZERS

Let us start this section with proving some auxiliary statements about free metabelian Lie algebras. We need them to describe annihilators and centralizers of partially commutative Lie algebras.

Lemma 4.1. *Suppose $u = [x_{j_1}, x_{j_2}, \dots, x_{j_m}] \in \mathfrak{B}(X)$, where $m \geq 2$. Let v be the largest monomial in the representation of $u.z$ as a linear combination of basis elements of $M(X)$. Then the first letter of v is x_{j_1} and the coefficient by this monomial is equal to 1.*

Proof. Let us consider the representation of $u.z$ as a linear combination of the elements of the form (2). We need to consider two cases.

1. Let $z \geq x_{j_2}$. In this case,

$$\begin{aligned} v &= [x_{j_1}, x_{j_2}, x_{j_3}, \dots, x_{j_m}].z \\ &= [x_{j_1}, x_{j_2}, \dots, x_{j_l}, z, x_{j_{l+1}}, \dots, x_{j_m}], \end{aligned}$$

where $x_{j_l} \leq z < x_{j_{l+1}}$. We obtain a monomial of the form (2) whose first letter is x_{j_1} .

2. Let $z < x_{j_2}$. Then

$$\begin{aligned} v &= [x_{j_1}, x_{j_2}, x_{j_3}, \dots, x_{j_m}].z \\ &= [x_{j_1}, x_{j_2}, z, x_{j_3}, \dots, x_{j_m}] \\ &= [x_{j_1}, z, x_{j_2}, x_{j_3}, \dots, x_{j_m}] + [\dots, [[x_{j_1}, [x_{j_2}, z]], x_{j_3}], \dots, x_{j_m}] \\ &= [x_{j_1}, z, x_{j_2}, x_{j_3}, \dots, x_{j_m}] - [x_{j_2}, z, x_{j_1}, x_{j_3}, \dots, x_{j_m}] \end{aligned}$$

We obtain a linear combination of two monomials of the form (2). Since x_{j_2} is the smallest letter of u the monomial with the first letter x_{j_1} is larger than the other one and the coefficient by this monomial is equal to 1. $\square$

Lemma 4.2. *Let $c = \alpha \bar{c} + \Delta c$ be the representation of $c \in M'(X)$ as a linear combination of basis elements with non-zero coefficients and let $\bar{c} = [y, x, x_{i_3}, \dots, x_{i_m}]$. Then for any $z \in X$ the largest monomial in the decomposition of $c.z$ as a linear combination of basis elements begins with the letter y and the coefficient by this monomial is equal to α .*

Proof. Let $c = \alpha\bar{c} + \Delta c$ be a homogeneous Lie monomial consisting of monomials of the form (2) and d a monomial appearing in Δc . Since $\bar{c}$ is the greatest monomial of c we have either $\text{mdeg}(\bar{c}) > \text{mdeg}(d)$ or $\text{mdeg}(\bar{c}) = \text{mdeg}(d)$ but $\bar{c}$ is greater than d lexicographically. Let us represent $\alpha(\bar{c}.z)$ and $(\Delta c).z$ as sums of elements of the form (2). By Lemma 4.1, the largest monomial of the decomposition of $\alpha(\bar{c}.z)$ begins with y and the coefficient by this monomial is α . On the other hand, the monomials in the decomposition of $(\Delta c).z$ are less than y with respect to the standard order. Let b be the largest monomial in the decomposition of $c.z$ to a linear combination of elements of the form (2). We have b begins with y and the coefficient by this monomial is equal to α . $\square$

The following result has been proved in [3] in the case of a free metabelian Lie algebra under a field. In his paper, we give the proof of this statement for a free metabelian Lie algebra over an arbitrary integral domain.

Theorem 4.3. *The derived subalgebra $M'(X)$ of $M(X)$ is a torsion-free $R[X]$ -module.*

Proof. We need to prove that $c.f \neq 0$ whenever $c \in M'(X) \setminus \{0\}$ and $f \in R[X] \setminus \{0\}$.

Note that $u.a \neq 0$ implies

$$(6) \quad \text{mdeg}(u.a) = \text{mdeg}(u) + \text{mdeg}(a)$$

for any Lie monomial u and any associative monomial a . So, without loss of generality it can be assumed that the decomposition of c to a linear combination of monomials of the form (2) is homogeneous by all generators and f is a monomial. Indeed, we can write $c = c_{\bar{\gamma}} + \sum_{\bar{\delta}} c_{\bar{\delta}}$, where $c_{\bar{\delta}}$ is a homogeneous Lie polynomial with the multidegree $\bar{\delta}$ and $\bar{\gamma}$ is the largest multidegree of the monomials in the decomposition of c . Let also $f = \alpha\bar{f} + \Delta f$, where $\bar{f}$ is the monomial of the largest degree appearing in f . It is sufficient to show that $c_{\bar{\gamma}}.\bar{f} \neq 0$. Indeed, by (6), the multidegrees of all other summands in the decomposition of $c.f$ are less than the multidegree of $c_{\bar{\gamma}}.\bar{f}$. Therefore, this monomial cannot cancel with the others.

So, suppose c is represented as a linear combination of monomials of the form (2), $\bar{c} = [y, x, x_{i_3}, \dots, x_{i_m}]$ is the largest monomial in this linear combination, and f is an associative monomial of the multidegree $\bar{\varepsilon}$. By $|\bar{\varepsilon}|$ denote the sum of coordinates of $\bar{\varepsilon}$. Let us proceed by induction on $|\bar{\varepsilon}|$.

If $|\bar{\varepsilon}| = 1$ then $f = z$ for some $z \in X$. Let b be the largest monomial in the representation of $c_{\bar{\gamma}}.z$ as a linear combination of elements of the form (2). By Lemma 4.2, b begins with y . The coefficient by this monomial is equal to the coefficient by $\bar{c}$. In particular, this coefficient is not equal to zero. Therefore, $c_{\bar{\gamma}}.z \neq 0$. By (6), this monomial cannot cancel with monomials in the decompositions of $c_{\bar{\delta}}.z$. Consequently, $c.z \neq 0$.

Suppose that the statement holds for all $\bar{\varepsilon}$ such that $|\bar{\varepsilon}| < r$. Let $|\bar{\varepsilon}| = r > 0$ and let f be a monomial of the multidegree $\bar{\varepsilon}$. For some generator x_i , we can write $f = \beta x_i f_0$, where $\beta \in R \setminus \{0\}$ and $|\text{mdeg} f_0| = r - 1$. We have $c.f = c.(\beta x_i f_0) = \beta(c.x_i).(f_0)$. As above, we get $(c.x_i) \neq 0$. Consequently, by the inductive hypothesis we obtain $(c.x_i).(f_0) \neq 0$. The proof is complete. $\square$

The following statement on commutative associative polynomials seems to be well-known, but anyway, we give its proof here.

Lemma 4.4. *Let R be an infinite integral domain and let $f \in R[x_1, \dots, x_n]$ be a non-zero polynomial over R in a finite number of indeterminates. Then there exist $r_1, \dots, r_n \in R$ such that $f(r_1, \dots, r_n) \neq 0$.*

Proof. Proceed by induction on the number of indeterminates. Let $f(x) = \alpha_l x^l + \dots + \alpha_1 x + \alpha_0$ be a non-zero polynomial in one indeterminate. Let us show that

this polynomial cannot have more than l roots. If y_1 is a root then there is the representation $f(x) = (x - y_1)f_1(x)$, where the degree of $f_1(x)$ is equal to $l - 1$. Similarly, one can find a root of $f_1(x)$, denote this root by y_2 , and obtain a similar decomposition for $f_1(x)$ and so on. Finally, we obtain the representation

$$(7) \quad f(x) = (x - y_1) \dots (x - y_s)g(x),$$

where $g(x)$ has no roots. Obviously, $f(x)$ has no roots except $y_1, \dots, y_s$. Indeed, if z is a root then substitution of z for x in (7) gives $f(z) = (z - y_1) \dots (z - y_s)g(z)$. Since R is an integral domain, at least one of the multiples is equal to 0 and $g(z) \neq 0$ because $g(x)$ has no roots. Therefore, $z = y_i$ for some i such that $1 \leq i \leq s$.

Since R is infinite, one can choose r not equal to any root of $f(x)$. Then $f(r) \neq 0$.

Suppose that the statement of the lemma holds for all polynomials in less than k indeterminates. Consider a polynomial f in k indeterminates. We have

$$(8) \quad f = \sum_{i=0}^l x_k^i f_i,$$

where $f_i \in R[x_1, \dots, x_{k-1}]$. Since $f \neq 0$ there exists i such that $f_i \neq 0$. By the inductive assumption, there are $r_1, \dots, r_{k-1}$ such that $f_i(r_1, \dots, r_{k-1}) \neq 0$ for some i . Substituting these values for the corresponding indeterminates in (8) we get a non-zero polynomial in the indeterminate x_k only. Consequently, there exists $r_k \in R$ such that substituting this element for x_k we obtain a non-zero element of R . So, the substitution $r_1, \dots, r_k$ for the corresponding indeterminates in (8) gives a non-zero element of R . This concludes the proof. $\square$

Lemma 4.5. *Let G be a graph such that each its connected component is a complete graph and $M(X; G)$ the partially commutative Lie algebra with the defining graph G . If $[x, y].f = 0$ for some vertices x and y and for some polynomial $f \in R[X]$ then either $[x, y] = 0$ (i.e. x and y belong to the same connected component of G) or $f = 0$.*

Proof. If G is a connected graph then $M(X; G)$ is just a free abelian Lie algebra. Therefore, $[x, y] = 0$ for any $x, y \in X$.

If G has at least two connected components then we proceed by induction on the number of generators in $M(X; G)$. Suppose $|X| = 2$ and G has two connected components (it means that G is totally disconnected). Then we have $M(X; G)$ is the free metabelian Lie algebra with two generators and the statement follows from Theorem 4.3.

Let the statement hold for all graphs having less than k vertices and satisfying the conditions of the lemma. Consider a graph G such that $|X| = k$. If this graph is totally disconnected then $M(X; G)$ is a free metabelian algebra and the statement follows from Theorem 4.3.

Suppose that G contains some edges and $[x, y].f = 0$, where $f \in R[X]$ and x and y are in different connected components. Consider an arbitrary connected component Γ of G such that Γ has at least two vertices. Let $Z = \{z, z_1, \dots, z_m\}$ be the set of all vertices in this component. If $x \in \Gamma$ (or $y \in \Gamma$) then let us set $z = x$ ($z = y$, respectively). By Y denote the set obtained from X by deleting vertices $z_1, \dots, z_m$.

Let $\varphi : M(X; G) \rightarrow M(Y; G_Y)$ be an identification on the connected component Γ defined as follows:

$$(9) \quad \varphi(x) = \begin{cases} \alpha_i z & \text{if } x = z_i \\ x & \text{else} \end{cases}.$$

We have the following representation of f :

$$f = \sum_j \gamma_j f_j z^{a_j} z_1^{a_{j,1}} \dots z_m^{a_{j,m}},$$

where f_j are monomials in $R[X \setminus Z]$. We obtain

$$\varphi(f) = \sum_j \gamma_j \alpha_1^{a_{j,1}} \dots \alpha_m^{a_{j,m}} f_j z^{a_j + a_{j,1} + \dots + a_{j,m}}.$$

Combining like terms by degrees of the elements in Y we obtain a sum such that each its coefficient is a polynomial in $\alpha_1, \dots, \alpha_m$. Moreover, since $f \neq 0$ at least one of the obtained coefficients is not equal to zero identically. Therefore, by Lemma 4.4, one can find $\alpha_1, \dots, \alpha_m$ such that $\varphi(f) \neq 0$. From (9), it follows that φ cannot map vertices from different connected components to one component. Consequently, $\varphi([x, y]) = [x, y] \neq 0$. Note that $M(Y; G_Y)$ satisfies the conditions of the lemma and the number of vertices in G_Y is less than k . Therefore, by the inductive hypothesis, $[x, y] \cdot \varphi(f) \neq 0$ in $M(Y; G_Y)$. It means that $[x, y] \cdot f \neq 0$ in $M(X; G)$. The proof is complete. $\square$

Definition 4.6. Let A be an associative commutative ring and let M be an A -module. An *annihilator* of a non-zero element $m \in M$ is the ideal A consisting of all elements $a \in A$ such that $m \cdot a = 0$.

Since for any partially commutative algebra $M(X; G)$ its derived subalgebra $M'(X; G)$ is an $R[X]$ -module, the annihilator of any element $g \in M'(X; G)$ in $R[X]$ is defined.

Let us move on to the description of annihilators in $M'(X; G)$. Define the ideal $I_{i,j}^G$ of $R[X]$ as follows. If x_i and x_j belong to different connected components in G then put $I_{i,j}^G = 0$. Let x_i and x_j be in the same connected component. For each path $(x_i, y_1, y_2, \dots, y_s, x_j)$ connecting these vertices in G consider the monomial $y_1 y_2 \dots y_s$. Define $I_{i,j}^G$ as the ideal generated by all such monomials. Obviously, it is sufficient to consider the monomials corresponding to the simple paths only.

Theorem 4.7. Let $M(X; G)$ be the partially commutative metabelian Lie algebra with the set of generators X and the defining graph G and let $x_i, x_j \in X$. If x_i and x_j are not adjacent then the annihilator of $[x_i, x_j]$ is equal to $I_{i,j}^G$.

Proof. First of all, let us show that if $f \in I_{i,j}^G$, then $[x_i, x_j] \cdot f = 0$. It is sufficient to prove this statement for the elements generating $I_{i,j}^G$.

Let us show that the statement is true for the paths having one intermediate vertex, i.e in the case $s = 1$. We have $[x_i, x_j] \cdot y_1 = [[x_i, x_j], y_1] = [[x_i, y_1], x_j] + [x_i, [x_j, y_1]] = 0$.

Let the statement hold for all simple paths having $s < k$ intermediate vertices. Suppose that the path $(x_i, y_1, \dots, y_k, x_j)$ has k intermediate vertices. We have

$$(10) \quad [x_i, x_j] \cdot (y_1 y_2 \dots y_k) = \sum [x_i \cdot (y_{i_1} \dots y_{i_t}), x_j \cdot (y_{j_1} \dots y_{j_{k-t}})],$$

where the sum is taken by all ordered partitions of the set $\{y_1, y_2, \dots, y_k\}$ by two subsets $\{y_{i_1} \dots y_{i_t}\}$ and $\{y_{j_1} \dots y_{j_{k-t}}\}$. Since $M(X; G)$ is metabelian it follows from (10) that:

$$\begin{aligned} [x_i, x_j] \cdot (y_1 y_2 \dots y_k) &= [x_i \cdot (y_1 \dots y_k), x_j] + [x_i, x_j \cdot (y_1 \dots y_k)] \\ &= [[x_i, y_1] \cdot (y_2 \dots y_k), x_j] + [x_i, [x_j, y_1] \cdot (y_2 \dots y_k)]. \end{aligned}$$

Since x_i are y_1 adjacent in G the first summand is equal to 0. The second summand contains the multiple $[x_j, y_1] \cdot (y_2 \dots y_k)$. Considering the path $(y_1, y_2, \dots, x_j)$ we obtain that this multiple is equal to zero by the induction hypothesis. Consequently, $[x_i, x_j] \cdot (y_1 y_2 \dots y_k) = 0$.

So, we have shown that $I_{i,j}^G$ is a subset of the annihilator of $[x_i, x_j]$. Let us prove the inverse statement. Without loss of generality, the annihilator of $[x_1, x_2]$ can be considered. There are two cases:

1. Suppose that x_1 and x_2 belong to different connected components of G . Let us add edges to the graph G in such a way that any two vertices belonging to the same connected component of G would be connected. Denote this graph by $\overline{G}$. So, connected components of G and $\overline{G}$ consist of the same vertices and each connected component of $\overline{G}$ is a complete graph.

Consider the identical simplification $\varphi : M(X, G) \rightarrow M(X, \overline{G})$. If $[x_1, x_2].f = 0$ in $M(X, G)$, then $\varphi([x_1, x_2].f) = 0$. Therefore, $[x_1, x_2].f = 0$ in $M(X, \overline{G})$. So, by Lemma 4.5, we have $f = 0$.

2. If x_1 and x_2 are not adjacent and belong to the same connected component, then we proceed by induction on $|X|$. Let $|X| = 3$. Then G has exactly two edges, namely $\{x_1, x_3\}$ and $\{x_2, x_3\}$. Indeed, otherwise, either x_1 and x_2 are adjacent or they belong to different connected components.

Consider the projection $\pi_Y : M(X; G) \rightarrow M(Y; G_Y)$, where $Y = \{x_1, x_2\}$. Let $[x_1, x_2].f = 0$. We have $\pi_Y([x_1, x_2].f) = [x_1, x_2].\pi_Y(f) = 0$. Since G_Y is totally disconnected $M(Y; G_Y) = M(Y)$. Therefore, by Theorem 4.3, $\pi_Y(f) = 0$. Consequently, $f = x_3 \hat{f}$.

Suppose that the statement holds for $|X| < n$. Consider an algebra $M(X; G)$, where X contains n vertices. Let $(x_1, y_1, y_2, \dots, y_s, x_2)$ be a simple path connecting x_1 and x_2 .

By X_i denote the set $X \setminus \{y_i\}$, where $i = 1, 2, \dots, s$. Let G_i be the graph generated by X_i .

Consider the set of projections $\pi_{X_i} : M(X; G) \rightarrow M(X_i; G_i)$, where $i = 1, 2, \dots, s$. Note that $\pi_{X_i}(x_1) = x_1$ and $\pi_{X_i}(x_2) = x_2$ for all i because the path $(x_1, y_1, y_2, \dots, y_s, x_2)$ is simple. Therefore, if $[x_1, x_2].f = 0$ in $M(X; G)$ then $\pi_{X_1}([x_1, x_2].f) = [x_1, x_2].\pi_{X_1}(f) = 0$ in $M(X_1; G_1)$. Let us remark that $M(X_1; G_1)$ is a subalgebra of $M(X; G)$ and $|X_1| = n - 1$. Consequently, by the inductive hypothesis, $\pi_{X_1}(f)$ is in the ideal $I_{1,2}^{G_1}$ of the algebra $M(X_1; G_1)$. In particular, $\pi_{X_1}(f)$ does not contain y_1 . Since π_{X_1} acts identically on all generators but y_1 there is the representation $f = \pi_{X_1}(f) + f_1$, where $f_1 \in R[X]$ is such that $\pi_{X_1}(f_1) = 0$. Consequently, $f_1 = y_1 \hat{f}_1$. It is clear that $I_{1,2}^{G_1} \subseteq I_{1,2}^G$. Therefore, $\pi_{X_1}(f) \in I_{1,2}^G$.

Suppose that for some $i < s$ there is the representation

$$(11) \quad f = f_i + \sum_{j=1}^i \tilde{f}_j,$$

where $\tilde{f}_j \in I_{1,2}^{G_j}$, and $f_i = y_1 y_2 \dots y_i \hat{f}_i$. Obviously, $I_{1,2}^{G_j} \subseteq I_{1,2}^G$ for $j = 1, 2, \dots, i$. Thus, we have $0 = [x_1, x_2].f = [x_1, x_2].f_i + \sum_{j=1}^i [x_1, x_2].\tilde{f}_j = [x_1, x_2].f_i$. Therefore, $0 = \varphi_{i+1}([x_1, x_2].f_i) = [x_1, x_2].\pi_{X_{i+1}}(f_i)$. So, by the induction hypothesis, $\pi_{X_{i+1}}(f_i)$ belongs to the ideal $I_{1,2}^{G_{i+1}}$ of the algebra $M(X_{i+1}; G_{i+1})$. As above, we obtain $f_i = f_{i+1} + \tilde{f}_{i+1}$, where $\tilde{f}_{i+1} = \pi_{X_{i+1}}(f_i) \in I_{1,2}^{G_{i+1}} \subseteq I_{1,2}^G$ and f_{i+1} is such that $\pi_{X_{i+1}}(f_{i+1}) = 0$. Therefore, $f_{i+1} = y_{i+1} \hat{f}_{i+1}$. Since each monomial of f_i is of the form $g = y_1 y_2 \dots y_i g'$ we obtain $f_{i+1} = y_1 y_2 \dots y_{i+1} \hat{f}_{i+1}$. Consequently, there is the following decomposition: $f = f_{i+1} + \sum_{j=1}^{i+1} \tilde{f}_j$. So, decomposition (11) holds for any i .

Setting $i = s$ in (11) we obtain $f = y_1 y_2 \dots y_s \widehat{f_s} + \sum_{j=1}^s \widetilde{f_j}$. As above, $\sum_{j=1}^s \widetilde{f_j} \in I_{1,2}^G$. Since $y_1 y_2 \dots y_s$ is a generating element of $I_{1,2}^G$, the monomial $y_1 y_2 \dots y_s \widehat{f_s}$ is also in this ideal. So, the proof is complete. $\square$

Before we move on to a description of centralizers in partially commutative metabelian Lie algebras we prove one more auxiliary assertion.

Let $X_{c,z} = X_c \cup \{z\}$ for any $z \in X \setminus X_c$, and let $G_{c,z}$ be the subgraph of G generated by $X_{c,z}$.

Remind that z is called a *cutpoint* of G if its deletion increases the number of connected components.

Lemma 4.8. *Let $c \in M'(X, G)$ be a non-zero element that can be written as a homogeneous linear combination of basis elements. If $c.z = 0$ for some $z \in X$ then z does not appear in c and it is a cutpoint of $G_{c,z}$.*

Proof. Let us reorder vertices of X in such a way that z would be not less than the vertices appearing in c . If c contains z or if z is greater than the smallest letter appearing in c then we can use the initial order on X . Otherwise, we can just permute z and the smallest letter of c in the initial linear order of X . Note that if c admits a representation as a homogeneous linear combination of basis elements under some order on X then it can be rewritten in such form under any other order on X . Moreover, for all orders on X the multidegrees of the monomials in the corresponding representations are same.

By Lemma 3.2, G_c has at least two connected components. Suppose that $\Gamma_0, \Gamma_2, \dots, \Gamma_s$ are connected components of G_c ($s \geq 1$). Let $y_0, y_1, \dots, y_s$ be the largest vertices of these components such that y_i lies in Γ_i for each $i = 0, \dots, s$. We may assume that $y_0 < y_1 < y_2 < \dots < y_s$. By x denote the minimal vertex appearing in c . Let x belong to Γ_0 .

We have:

$$(12) \quad c = \sum_{i=1}^s \beta_i [y_i, x, x_{i_3}, \dots, x_{i_m}],$$

where all summands have the same multidegree and some β_i are not equal to 0. Let $r \leq s$ be the largest number such that $\beta_r \neq 0$. Consider $c.z$ as an element of $M(X)$. Since $z \geq x$, the proofs of Lemma 4.1 (case 1) and Lemma 4.2 imply that the largest monomial of $c.z$ can be written as follows: $\overline{c}.z = \overline{c}.z = [y_r, x, x'_{r_3}, \dots, x'_{r_{m+1}}]$.

Let $c.z = 0$ in $M(X; G)$. If $\overline{c}.z = 0$ in $M(X; G)$, then Lemma 3.2 implies that y_r and x belong to the same connected component of $G_{c,z}$, but in different connected components of G_c . Consequently, z does not appear in c and there are vertices in Γ_0 and Γ_r adjacent to z . It means that z is a cutpoint of $G_{c,z}$.

If $\overline{c}.z \neq 0$ in $M(X; G)$, then the monomial $\overline{c}.z$ is equal to $c' = [y_j, x, \dots, x'_{j_{m+1}}]$ for some $j \neq r$ in this algebra and $\text{mdeg}(\overline{c}) = \text{mdeg}(c')$. But it is possible only if y_j and y_r are in the same connected component of $G_{c,z}$. Therefore, z does not appear in c and it is a cutpoint of $G_{c,z}$. $\square$

Let G be an arbitrary graph with the set of vertices X and let u be an arbitrary element of $M(X; G)$. By $C_G(u)$ denote the centralizer of $u \in X$ in the algebra $M(X; G)$. So, $C_G(u)$ is the set of all elements $v \in M(X; G)$ such that $[v, u] = 0$. Let us also introduce the notation $\mathcal{C}_G(x)$ for $C_G(x) \cap M'(X; G)$. We are going to describe the centralizers of the elements in X and their linear combinations. We need this description in the proof of Theorem 5.9 in Sec. 5. The following theorem holds.

Theorem 4.9. *Let G be a graph with the set of vertices $X = \{x_1, x_2, \dots, x_n\}$.*

1. *If x_n is an isolated vertex in G then $C_G(x_n)$ consists of the elements v of the*

form

$$(13) \quad v = \alpha_n x_n,$$

where $\alpha_n \in R$.

2. If the degree of x_n is equal to 1 in G (say, it is adjacent to x_{n-1}) then $C_G(x_n)$ consists of all elements v of the form

$$(14) \quad v = \alpha_{n-1} x_{n-1} + \alpha_n x_n,$$

where $\alpha_{n-1}, \alpha_n \in R$.

3. If x_n is adjacent to $x_{r+1}, \dots, x_{n-1}$ in G ($r \leq n-3$), then $C_G(x_n)$ consists of all elements v of the form

$$(15) \quad v = \sum_{k=r+1}^n \alpha_k x_k + \sum_{r+1 \leq i < j \leq n-1} [x_i, x_j] \cdot f_{ij},$$

where $\alpha_k \in R$, $f_{ij} \in R[X \setminus \{x_n\}]$.

Proof. Clearly, all elements of the form (13) or (14) (depending on the number of vertices adjacent to x_n) in $M(X; G)$ are in $C_G(x_n)$. It is also easy to see that if x_n is adjacent to $x_{r+1}, \dots, x_{n-1}$ then any element of the form $v = \sum_{k=r+1}^n \alpha_k x_k$ belongs to the centralizer of x_n . So, we are left to show that the elements of the form $\sum_{r+1 \leq i < j \leq n-1} [x_i, x_j] \cdot f_{ij}$ (see case 3) are in $C_G(x_n)$. We need to prove that

$$\sum_{r+1 \leq i < j \leq n-1} [x_i, x_j] \cdot f_{ij} x_n = 0.$$

It follows from Lemma 3.2. Indeed, let us consider the element of the form $w = [x_i, x_j] \cdot f_{ij} x_n$. Since x_i and x_j are adjacent to x_n , the vertices x_i and x_j are in the same connected component of G_w and we are done.

Conversely, suppose that $c \in C_G(x_n)$. Let us write this element as a linear combination of basis monomials. As we noticed in Sec. 2 if a Lie polynomial is equal to zero in $M(X; G)$ then all homogeneous components (in the sense of multidegree) of this polynomial are equal to zero in $M(X; G)$. So, without loss of generality it can be assumed that all basis monomials of c have the same multidegree. Consider two cases.

1. If a homogeneous element $c \in C_G(x_n)$ is a linear combination of monomials of the length 1, then $c = \alpha_i x_i$ for some i . Consequently, $0 = [c, x_n] = \alpha_i [x_i, x_n]$. So, it is easy to see that either $i = n$ or x_i is adjacent to x_n .

2. Suppose that $c \in C_G(x_n)$. Let us represent c as a linear combination of basis elements of the algebra $M(X; G)$. Consider the graph G_c . Let this graph have $s+1$ connected components ($s \geq 1$). By $y_0, y_1, y_2, \dots, y_s$ denote the largest vertices of the connected components as we did in the proof of Lemma 4.8. Let x be the smallest letter appearing in c . We can suppose that x is in the same connected component as y_0 . Since $[c, x_n] = 0$, Lemma 4.8 implies, that x_n is a cutpoint of G_{c, x_n} .

Clearly, degree of x_n in G is not less than 2. Indeed, x_n should be connected with at least two connected components of G_c . Therefore, its degree in G_{c, x_n} should be greater than 1. So, if the degree of x_n in G is not greater than 1 then $C(x_n) = 0$ and parts 1 and 2 hold.

Thus, x_n is a cutpoint of G_{c, x_n} . We can assume without loss of generality that one of the connected components contains the vertices $y_1, \dots, y_t$ for $t \leq s$, and perhaps y_0 . Moreover, one can suppose that $y_1, y_2, \dots, y_t$ are adjacent to x_n , i.e.

$$(16) \quad \{y_1, \dots, y_t\} \subseteq \{x_{r+1}, \dots, x_n\}.$$

We can also assume that if y_0 is in the same connected component of G_{c,x_n} as the vertices $y_1, \dots, y_t$ then y_0 is also adjacent to x .

Since the representation of c in the form (12) holds we obtain

$$(17) \quad [c, x_n] = \sum_{i=1}^s \beta_i [y_i, x, x_{i_3}, \dots, x_{i_m}, x_n] \\ = \sum_{i=1}^t \beta_i [x_n, x, x_{i_3}, \dots, x_{i_m}, y_i] + \sum_{j=t+1}^s \beta_j [y_j, x, x_{j_3}, \dots, x_{j_m}, x_n].$$

By Theorem 3.3, the monomials $[y_j, x, x_{j_3}, \dots, x_{j_m}, x_n]$ are linearly independent for $t+1 \leq j \leq s$. By the same theorem, we see that the first sum of the right hand side (17) is equal to a multiple of a monomial of the corresponding multidegree beginning with x_n which is either equal to zero in $M(X; G)$ or a basis element. In the latter case, this element is linearly independent from the elements of the second sum. Therefore, in both cases $\beta_j = 0$ if $t+1 \leq j \leq s$. By (12) we obtain

$$(18) \quad c = \sum_{i=1}^t \beta_i [y_i, x, x_{i_3}, \dots, x_{i_m}].$$

Consequently,

$$(19) \quad [c, x_n] = \sum_{i=1}^t \beta_i [y_i, x, x_{i_3}, \dots, x_{i_m}, x_n].$$

In the former case, we obtain x_n is in the same connected component of G_{c,x_n} as x and y_0 . In this case, by Lemma 3.2 we have $[y_i, x, x_{i_3}, \dots, x_{i_m}, x_n] = 0$ for $i \leq t$. On the other hand, Lemma 3.1 implies that $[y_i, x, x_{i_3}, \dots, x_{i_m}, x_n] = [y_i, y_0, x'_{j_3}, \dots, x'_{i_m}, x'_{i_{m+1}}]$ and y_0 and y_i are adjacent to x_n . So, as follows from (16), c can be represented in the form (15).

In the latter case, we have y_0 and x_n are in different connected components of G_{c,x_n} . By (19), we obtain the following equation in $M(X; G)$:

$$0 = [c, x_n] = \sum_{i=1}^t \beta_i [x_n, x, x_{i_3}, \dots, x_{i_m}, y_i].$$

Let us arrange the last $m-1$ multiples in each summand of the right-hand side of this equation in the increasing order. We obtain a linear combination of t equal monomials which are not equal to zero in $M(X; G)$. Since $[c, x_n] = 0$ we obtain $\sum_{i=1}^t \beta_i = 0$. So, the following equation holds

$$(20) \quad c = \sum_{i=1}^{t-1} \beta_i ([y_i, x, x_{i_3}, \dots, x_{i_m}] - [y_t, x, x_{t_3}, \dots, x_{t_m}]).$$

For each summand of the right-hand side of (20), we can apply the Lie algebra identities and rearrange the last $m-2$ multiples. So, we obtain

$$(21) \quad [y_i, x, x_{i_3}, \dots, x_{i_m}] - [y_t, x, x_{t_3}, \dots, x_{t_m}] \\ = [y_i, x, y_t, x'_4, \dots, x'_m] - [y_t, x, y_i, x'_4, \dots, x'_m] \\ = [y_i, x, y_t, x'_4, \dots, x'_m] + [x, y_t, y_i, x'_4, \dots, x'_m] \\ = -[y_t, y_i, x, x'_4, \dots, x'_m] \\ = [y_i, y_t, x, x'_4, \dots, x'_m]$$

By (20) and (21), we get

$$c = \sum_{i=1}^{t-1} \beta_i [y_i, y_t, x, x'_4, \dots, x'_m].$$

Therefore, c can be represented in the form (15). This completes the proof of part 3. $\square$

Now we are ready to describe centralizers of linear combinations of the elements in X . The following lemma holds:

Lemma 4.10. *Let $c = c_0 + \Delta c$, where c_0 is a homogeneous component of c having the largest multidegree. If $c \in \mathcal{C}(\sum_{j=1}^l \alpha_j x_{i_j})$, where all α_j are not equal to 0, then $c_0 \in \bigcap_{j=1}^l \mathcal{C}(x_{i_j})$.*

Proof. To be definite assume that

$$c \in \mathcal{C}(\sum_{j=k+1}^n \alpha_j x_j).$$

So,

$$\begin{aligned} 0 &= c.(\sum_{j=k+1}^n \alpha_j x_j) \\ (22) \quad &= \sum_{j=k+1}^n \alpha_j [c, x_j] \\ &= \sum_{j=k+1}^{n-1} \alpha_j [c_0, x_j] + \alpha_n [c_0, x_n] + \sum_{j=k+1}^n \alpha_j [\Delta c, x_j]. \end{aligned}$$

Moreover, $[c_0, x_n] = 0$. Indeed, otherwise, the basis monomials appearing in the decomposition of $[c_0, x_n]$ have the greatest multidegree in the decomposition of $c.(\sum_{j=k+1}^n \alpha_j x_j)$. So, they cannot cancel with other summands. Therefore, $c_0 \in \mathcal{C}(x_n)$.

By Lemma 4.8, c_0 does not contain x_n in it, neither does c . Let $X' = X \setminus \{x_n\}$ and let G' be the subgraph of G generated by X' . Consider the projection $\pi_{X'}$. This projection maps c identically to $M(X'; G')$. Arguing as above, we see that $c_0 \in \mathcal{C}(x_{n-1})$, and so on. Finally, we get $c_0 \in \bigcap_{j=k+1}^n \mathcal{C}(x_j)$, that concludes the proof. $\square$

Theorem 4.11. *Let $M(X; G)$ be a partially commutative metabelian Lie algebra, where $X = \{x_1, x_2, \dots, x_n\}$. Then*

$$\mathcal{C}(\sum_{j=1}^m \alpha_{i_j} x_{i_j}) = \bigcap_{j=1}^m \mathcal{C}(x_{i_j})$$

for any elements $x_{i_1}, x_{i_2}, \dots, x_{i_m}$ and for any $\alpha_{i_1}, \alpha_{i_2}, \dots, \alpha_{i_m} \in R \setminus \{0\}$.

Proof. It is sufficient to show that $\mathcal{C}(\sum_{j=k+1}^n \alpha_j x_j)$, because otherwise, we can just renumber the vertices as we did in the proof of Lemma 4.10.

The inclusion

$$\bigcap_{j=1}^m \mathcal{C}(x_{i_j}) \subseteq \mathcal{C}(\sum_{j=1}^m \alpha_{i_j} x_{i_j})$$

is obvious. So, we are left to show that

$$\mathcal{C}(\sum_{j=k+1}^n \alpha_j x_j) \subseteq \bigcap_{j=k+1}^n \mathcal{C}(x_j).$$

Let $c \in \mathcal{C}(\sum_{j=k+1}^n \alpha_j x_j)$. Represent c as a linear combination of elements in $\mathfrak{B}(X; G)$. We obtain $c = c_0 + \Delta c$, where c_0 is a homogeneous component of c having the largest multidegree. By Lemma 4.10, $c_0 \in \bigcap_{j=k+1}^n \mathcal{C}(x_j)$. So, we obviously have $c_0 \in \mathcal{C}(\sum_{j=k+1}^n \alpha_j x_j)$. Consequently, $\Delta c = c - c_0 \in \mathcal{C}(\sum_{j=k+1}^n \alpha_j x_j)$ and the decomposition of Δc to the sum of homogeneous components has one summand less than the similar decomposition of c . We can apply Lemma 4.10 for Δc and so on. After finitely many steps we obtain each homogeneous component of c belongs to $\bigcap_{j=k+1}^n \mathcal{C}(x_j)$, so does their sum. Since this sum is equal to c we are done. $\square$

In Sec. 5, we consider partially commutative Lie algebras whose defining graphs are trees. Let us prove the following theorem about centralizers in this case.

Theorem 4.12. *Let G be a tree and let $\alpha_j \in R \setminus \{0\}$ for $j = 1, 2, \dots, m$, where $m \geq 2$. Then $\mathcal{C}(\sum_{j=1}^m \alpha_j x_{i_j}) = 0$.*

Proof. Since $\mathcal{C}(\sum_{j=1}^m \alpha_j x_{i_j}) \subseteq \mathcal{C}(\sum_{j=1}^{m-1} \alpha_j x_{i_j})$, it is sufficient to show that $\mathcal{C}(\alpha x_s + \beta x_t) = 0$, if $\alpha, \beta \neq 0$ and $s \neq t$.

If either x_s or x_t is an endpoint, then Theorem 4.9 implies that the corresponding centralizer is trivial and the statement holds obviously. So, we are left to consider the case when neither vertex is an endpoints.

Let $c \in \mathcal{C}(\alpha x_s + \beta x_t)$. By the proof of Theorem 4.11, we may assume that c is a homogeneous element of $M(X; G)$. By Lemma 4.10, neither x_s , nor x_t appears in c . Therefore, these vertices are not in G_c .

By $\tilde{X}$ denote the set $X_c \cup \{x_s, x_t\}$. Let $\tilde{G}$ be the subgraph of G generated by $\tilde{X}$. It is obvious that $\tilde{G}$ is a forest.

Consider the graph G_c . This graph is also a forest. Moreover, all vertices adjacent to x_s are in different connected components of G_c . Indeed, let y and z be adjacent to x_s and belong to the same connected component of G_c . Then there is a simple path $(y, w_1, w_2, \dots, w_l, z)$ in this graph. However, since x_s is not in G_c we obtain the cycle $(x_s, y, w_1, w_2, \dots, w_l, z, x_s)$ in G . This is a contradiction because G is a tree. It is obvious that the assertion also holds for the vertices adjacent to x_t .

Let us show that there is at most one connected component Γ of G_c satisfying the following property: there are vertices $z, z' \in \Gamma$ such that z is adjacent to x_s and z' is adjacent to x_t . Indeed, if there are at least two such components (denote two of them as Γ_1 and Γ_2), then there exists a cycle going from x_s to x_t by vertices of Γ_1 and then, backward, from x_t to x_s by vertices of Γ_2 . Since G is a tree, we get a contradiction.

So, G_c has at most one connected component having a vertex adjacent to x_s as well as a vertex adjacent to x_t . If Γ exists, we can renumber the vertices of G in such a way that the minimal vertex (denote it by x) is in Γ and each vertex adjacent to either x_s or x_t and not belonging to Γ is the largest in its connected component. If there is no component Γ then any vertex of G_c can be chosen as x .

Theorem 4.11 implies that $c \in \mathcal{C}(x_s) \cap \mathcal{C}(x_t)$. By Theorem 4.9, we obtain

$$(23) \quad c = \sum_{\{x_i, x_s\}, \{x_j, x_s\} \in G_c} [x_i, x_j] \cdot f_{ij}$$

and

$$(24) \quad c = \sum_{\{x_p, x_t\}, \{x_q, x_t\} \in G_c} [x_p, x_q] \cdot g_{pq}$$

FIGURE 1. Subgraph generated by z_1, z_2, z_3, z_4

where $f_{ij}, g_{pq} \in R[X_c]$ are such that the multidegree of each monomial in both equations is equal to $\text{mdeg}(c)$. If $x = x_i$ in the monomial $[x_i, x_j] \cdot f_{ij}$ of the right-hand side of (23), then it can be written in the form

$$(25) \quad [x_i, x_j] \cdot f_{ij} = -[x_j, x] \cdot f_{ij}.$$

If $x \neq x_i, x_j$, then put the smallest letter in each monomial to the third position. After that, we can transform each such summand in the right-hand side of (23) as follows:

$$(26) \quad \begin{aligned} [x_i, x_j] \cdot f_{ij} &= \sum_l \alpha_l [x_i, x_j, x, y_1, \dots, y_m] \\ &= \sum_l \alpha_l ([x_i, x, x_j, y_1, \dots, y_m] - [x_j, x, x_i, y_1, \dots, y_m]) \end{aligned}$$

Substituting (25) and (26) to (23) and combining like terms we obtain a linear combination of basis monomials such that the first letter of each of them is adjacent to x_s and does not belong to Γ (if there is the component Γ in G_c). Indeed, if, for instance, x_i and x belong to the same connected component of G_c then $[x_i, x, x_j, y_1, \dots, y_m] = 0$. It means that this summand is not in the basis of $M(X; G)$ and we can exclude it from the obtained linear combination. Let us rewrite each summand in the right-hand side of (24) in a similar way. We obtain a linear combination of basis elements, each of which begins with a letter adjacent to x_t and not belonging to Γ . Since no vertices outside Γ can be adjacent to both x_s and x_t the obtained linear combinations have no equal monomials. Since the set of monomials in (23) and (24) is linearly independent, all coefficients by the basis elements in these linear combinations are equal to zero. Therefore, $c = 0$ and the proof is complete. $\square$

5. UNIVERSAL EQUIVALENCE

In this section, $M(X; T)$ denotes a partially commutative metabelian Lie ring (i.e. a partially commutative metabelian Lie algebra over the ring of integers $\mathbb{Z}$) whose defining graph is a tree.

As above, let $X = \{x_1, x_2, \dots, x_n\}$. Denote by x the smallest vertex in X (namely the vertex x_1).

We need some auxiliary technical results to prove the main theorem.

Lemma 5.1. *Let $T_1 = \langle X, E_1 \rangle$ and $T_2 = \langle Y, E_2 \rangle$ be trees. If the algebras $M(X; T_1)$ and $M(Y; T_2)$ are universally equivalent and at least two vertices of T_1 are not endpoints, then at least two vertices of T_2 are not endpoints either.*

Proof. Let $Y = \{y_1, y_2, \dots, y_m\}$ and set $y_i > y_j$ iff $i > j$. First of all, let us show that if at least two vertices of T_1 are not endpoints then at least two of them are adjacent. Suppose that x_i and x_j are not endpoints. If x_i and x_j are not adjacent in T_1 then there exists a simple path $(x_i, x_{i_1}, \dots, x_{i_k}, x_j)$ connecting these vertices. We are left to notice that all inner vertices of this path are not endpoints because the degree of each endpoint is 1 while the degree of any inner vertex is at least 2.

Therefore, we can choose different vertices z_1, z_2, z_3, z_4 in X in such a way that the subgraph $\tilde{T}$ of T_1 generated by these vertices seems as on Figure 1. Denote

$\{z_1, z_2, z_3, z_4\}$ by $\tilde{X}$.

Since $\tilde{T}$ has no cycles the following formula holds in $M(\tilde{X}; \tilde{T})$ and therefore, in $M(X; T_1)$:

$$(27) \quad \begin{aligned} \exists v_1 v_2 v_3 v_4 ([v_1, v_2] = [v_2, v_3] = [v_3, v_4] = 0 \\ \wedge [v_1, v_3] \neq 0 \wedge [v_1, v_4] \neq 0 \wedge [v_2, v_4] \neq 0). \end{aligned}$$

We can put v_i to be equal to z_i .

Suppose that T_2 has at most one vertex which is not an endpoint. If all vertices of T_2 are endpoints then Y contains one or two vertices. In this case $M(Y; T_2)$ is an abelian Lie algebra. Consequently, formula (27) does not hold in this algebra.

So, without loss of generality we may assume that y_m is not an endpoint. Then this vertex is adjacent to all other vertices and there are no other edges. So,

$$(28) \quad M(Y; T_2) = M(Y') \oplus \mathcal{L}(y_m),$$

where $Y' = Y \setminus \{y_m\}$, $\mathcal{L}(y_m)$ is one-dimensional Lie algebra, and $[u, v] = 0$ for any $u \in M(Y')$ and any $v \in \mathcal{L}(y_m)$. It means that $M(Y; T_2)$ is a direct sum of a free metabelian Lie algebra and the one-dimensional abelian Lie algebra (as vector spaces) and Lie product of any elements from different direct summands is equal to zero.

Suppose that $u_1, u_2, u_3 \in M(Y')$. Let us show that if $[u_1, u_2] = [u_2, u_3] = 0$ then $[u_1, u_3] = 0$.

If $u_2 \notin M'(Y')$ then Theorem 2.2 implies that there exist $\alpha, \beta, \gamma, \delta \in \mathbb{Z} \setminus \{0\}$, such that $\alpha u_1 = \beta u_2$, $\gamma u_2 = \delta u_3$. Consequently, $\alpha \gamma u_1 = \beta \gamma u_2 = \beta \delta u_3$, where $\alpha \gamma, \beta \delta \neq 0$. It means that u_1 and u_3 are linearly dependent. Therefore, $[u_1, u_3] = 0$ by Theorem 2.2.

Again by Theorem 2.2, it follows that if $u_2 \in M'(Y')$ then $u_1, u_3 \in M'(Y')$. Therefore, $[u_1, u_3] = 0$.


So, if T_1 has at least two vertices which are not endpoints, then (27) holds in $M(X; T_1)$. Since $M(X; T_1)$ and $M(Y; T_2)$ are universally equivalent, formula (27) holds in $M(Y; T_2)$. By (28), each element v_i can be represented in the form $v_i = u_i + \alpha_i y_m$, where $u_i \in M(Y')$. Since the multiples of y_m are in the center of $M(Y, T_2)$, we obtain $[v_i, v_j] = [u_i, u_j]$.

Consequently, if $[v_1, v_2] = [v_2, v_3] = 0$ then $[u_1, u_2] = [u_2, u_3] = 0$ and we have $[u_1, u_3] = 0$. Therefore, $[v_1, v_3] = 0$ and we have a contradiction to (27). Thus, at least two vertices of T_2 are not endpoints vertices. $\square$

Lemma 5.2. *Let T be a tree and let X be the set of its vertices at least k of which are not endpoints. Then the following formula holds in $M(X; T)$:*

$$(29) \quad \begin{aligned} \Phi(T) = \exists z_1 \dots z_k u_1 \dots u_k v_1 \dots v_k \left(\bigwedge_{i=1}^k [u_i, v_i] \neq 0 \wedge \bigwedge_{i=1}^k [u_i, v_i, z_i] = 0 \wedge \right. \\ \left. \bigwedge_{i \neq j} [u_i, v_i, z_j] \neq 0 \wedge \bigwedge_{\{x_i, x_j\} \in T} [z_i, z_j] = 0 \wedge \bigwedge_{\{x_i, x_j\} \notin T} [z_i, z_j] \neq 0 \right). \end{aligned}$$

Proof. Without loss of generality we may assume that $x_1, \dots, x_k$ are not endpoints. These vertices can be chosen for $z_1, \dots, z_k$ respectively. For each $i = 1, 2, \dots, k$, let us take two different vertices adjacent to x_i as u_i and v_i . We have $[u_i, v_i] \neq 0$. Indeed, if u_i is adjacent to v_i then (u_i, v_i, z_i, u_i) is a cycle in T and we obtain a contradiction. Next, $[u_i, v_i, z_i] = [u_i, z_i, v_i] + [u_i, [v_i, z_i]] = 0$ since z_i is adjacent to both u_i and v_i . On the other hand, if $i \neq j$ then $[u_i, v_i, z_j] \neq 0$. Indeed, otherwise Lemma 3.2 implies that u_i and v_i are in the same connected component of the subgraph of T generated by the set $\{u_i, v_i, z_j\}$. Consequently, either u_i is adjacent

FIGURE 2. Graph T

to v_i or the vertex z_j is adjacent to both u_i and v_i . The former case contradicts to the inequality $[u_i, v_i] \neq 0$. In the latter case, we have $(u_i, z_i, v_i, z_j, u_i)$ is a cycle. Since T is a tree we again have a contradiction. Finally, it is obvious that $[z_i, z_j] = 0$ for $\{x_i, x_j\} \in T$ and $[z_i, z_j] \neq 0$ for $\{x_i, x_j\} \notin T$. $\square$

Suppose that T has a vertex adjacent to at least two endpoints. Without loss of generality we can assume that x is such vertex, moreover, the vertices x_{n-2}, x_{n-1}, x_n are adjacent to x , and x_n is an endpoint (see Figure 2).

Denote by X' the set $X \setminus \{x_n\}$. Let, $T_{X'}$ be the subtree of T generated by X' .

For any $\lambda, p \in \mathbb{Z}$ we define the map $\varphi_{\lambda, p} : X \rightarrow M(X'; T_{X'})$ as follows:

$$\varphi_{\lambda, p}(x_i) = \begin{cases} x_i, & \text{if } i \neq n, \\ \lambda p x_{n-1} + \lambda x_{n-2}, & \text{if } i = n. \end{cases}$$

It is easy to see that this map can be extended up to a homomorphism from $M(X; T)$ to $M(X'; T_{X'})$ uniquely. We denote this homomorphism by $\varphi_{\lambda, p}$ as well. Let $\mathbb{Z}^+$ denote the set of all positive integers.

Lemma 5.3. *Let $f(x, y)$ be a non-zero polynomial over $\mathbb{R}$. Then there exist a number $X \in \mathbb{Z}^+$ and a function $Y : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ such that $f(x, y) \neq 0$ for all positive integers $x > X$ and $y > Y(x)$.*

Proof. Rewrite $f(x, y)$ in the form

$$f(x, y) = \sum_{i=0}^n p_i(x) y^i,$$

where $p_i(x)$, $i = 0, 1, \dots, n$ are polynomials in the indeterminate x . Let X be the smallest positive integer that is greater than all roots of all polynomials $p_i(x)$. Then for each $x_0 > X$ we have $g(y) = f(x_0, y)$ is a polynomial in y with non-zero coefficients. Consequently, one may set $Y(x_0)$ to be the smallest positive integer that is greater than the greatest root of $g(y)$. One can choose any values for $Y(x_0)$ if $x_0 \leq X$. $\square$

Lemma 5.4. *Let $g \in M(X; T) \setminus M'(X; T)$ and p any positive integer; then there exists $\lambda_0 \in \mathbb{Z}^+$ such that $\varphi_{\lambda, p}(g) \neq 0$ for any $\lambda \geq \lambda_0$.*

Proof. Let $g = g_0 + g_1$, where $g_0 = \sum_{i=1}^n \alpha_i x_i$ and $g_1 \in M'(X; G)$. Since $g \in M(X; G) \setminus M'(X; G)$ we have $g_0 \neq 0$.

If $\alpha_n = 0$, then $\varphi_{\lambda,p}(g) = g_0 + \varphi_{\lambda,p}(g_1)$. Since $\varphi_{\lambda,p}(g_1) \in M'(X; T)$ one can see that g_0 and $\varphi_{\lambda,p}(g_1)$ are linearly independent. Thus, $\varphi_{\lambda,p}(g) \neq 0$ for any λ and we can put, for example, $\lambda_0 = 1$.

If $\alpha_n \neq 0$, then

$$\varphi_{\lambda,p}(g) = \sum_{i=1}^{n-3} \alpha_i x_i + (\alpha_{n-2} + \lambda \alpha_n) x_{n-2} + (\alpha_{n-1} + \lambda p \alpha_n) x_{n-1} + \varphi_{\lambda,p}(g_1).$$

Since $\mathbb{Z}$ is an integral domain and the polynomial $\alpha_{n-2} + \lambda \alpha_n$ does not depend on p there exists λ_0 satisfying the conditions of the lemma. So, $(\alpha_{n-2} + \lambda \alpha_n) x_{n-2} \neq 0$. Since all generators are linearly independent, this summand cannot cancel with other summands of the length 1. Therefore, $\varphi_{\lambda,p}(g_0) \neq 0$. It means that $\varphi_{\lambda,p}(g_0) \notin M'(X; T)$. Since $\varphi_{\lambda,p}(g_1) \in M'(X; T)$, we obtain $\varphi_{\lambda,p}(g) = \varphi_{\lambda,p}(g_0) + \varphi_{\lambda,p}(g_1) \neq 0$ for any $\lambda \geq \lambda_0$. $\square$

Consider $g \in M'(X; T)$. There is the decomposition

$$(30) \quad g = \sum_{\bar{\delta}} g_{\bar{\delta}},$$

where each $g_{\bar{\delta}}$ is a Lie monomial such that $\text{mdeg}(g_{\bar{\delta}}) = \bar{\delta}$. Denote by $\text{mdeg}(v; k)$ the tuple of the first k coordinates of $\text{mdeg}(v)$. By analogy, $\bar{\delta}(k)$ is the tuple $(\delta_1, \delta_2, \dots, \delta_k)$.

Lemma 5.5. *Let $g \in M(X; T)$ and let there be a positive integer number N such that for any $\lambda > N$ the equality $\varphi_{\lambda,p}(g) = 0$ holds for infinitely many integers $p > P(\lambda)$. Then $\varphi_{\lambda,p}(g_{\bar{\delta}}) = 0$ for all λ, p and for all $g_{\bar{\delta}}$ in decomposition (30).*

Proof. Consider $g_{\bar{\delta}}$ for some $\bar{\delta}$. Let v be a basis monomial in the decomposition of $g_{\bar{\delta}}$ and v' a monomial in the decomposition of $\varphi_{\lambda,p}(v)$ as a linear combination of basis elements. Then $\text{mdeg}(v'; n-3) = \bar{\delta}(n-3)$.

Let v' be some monomial of $M(X'; T')$. Consider a basis monomial v of $M(X; T)$ such that v' is contained in the decomposition of $\varphi_{\lambda,p}(v)$ as a linear combination of basis elements. We have $\text{mdeg}(v; n-3) = \text{mdeg}(v'; n-3)$. Moreover, it is easy to see that $\text{mdeg}_{n-2}(v) + \text{mdeg}_{n-1}(v) + \text{mdeg}_n(v) = \text{mdeg}_{n-2}(v') + \text{mdeg}_{n-1}(v')$, $\text{mdeg}_{n-2}(v) \leq \text{mdeg}_{n-2}(v')$, and $\text{mdeg}_{n-1}(v) \leq \text{mdeg}_{n-1}(v')$. Hence, if $\text{mdeg}(v') = (\gamma_1, \dots, \gamma_{n-1})$ then $\text{mdeg}(v) = (\gamma_1, \dots, \gamma_{n-3}, \gamma_{n-2} - s, \gamma_{n-1} - r, r + s)$, where $r, s \geq 0$.

Note that the coefficient of v' in the decomposition of $\varphi_{\lambda,p}(v)$ is $\beta \lambda^{r+s} p^r$, where β does not depend on λ, p .

Let $g = \sum \alpha_i u_i$, where u_i are different basis monomials. Let us find $\varphi_{\lambda,p}(u_i)$ for all u_i and compute the sum of coefficients by v' in all such monomials. This sum is equal to

$$\sum_{m=0}^{\gamma_{n-2}+\gamma_{n-1}} \sum_{k=\max(0, m-\gamma_{n-2})}^{\min(m, \gamma_{n-1})} \beta_{m,k} \lambda^m p^k,$$

where the coefficients $\beta_{m,k}$ depend on the coefficients α_i but do not depend on λ and p .

On the other hand, since $\varphi_{\lambda,p}(g) = 0$ for $\lambda > N$ and for infinitely many positive integers p , we obtain the coefficient by v' is equal to zero. By Lemma 5.3, we get $\beta_{m,k} = 0$ for all m and k . We are left to note that $\beta_{m,k} \lambda^m p^k$ is the sum of coefficients by v' in the decomposition of the elements $\varphi_{\lambda,p}(u_i)$ such that $\text{mdeg}(u_i) = (\gamma_1, \dots, \gamma_{n-3}, \gamma_{n-2} - m + k, \gamma_{n-1} - k, m)$. So, the multidegrees of all such elements are same. Therefore, the coefficient by v' in the decomposition of $\varphi_{\lambda,p}(g_{\bar{\delta}})$ is 0 for any multidegree $\bar{\delta}$. Since v' is an arbitrary monomial, the proof is complete. $\square$

Let $w(x_i, \bar{\delta}(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n)$ be the basis monomial of $M(X; T)$ such that it starts with x_i and its multidegree is $\bar{\delta} = (\delta_1, \dots, \delta_n)$. Similarly, let us denote by $w'(x_i, \bar{\gamma}(n-3), \gamma_{n-2}, \gamma_{n-1})$ the basis monomial of $M(X'; T')$ such that it starts with x_i and its multidegree is $\bar{\gamma} = (\gamma_1, \dots, \gamma_{n-1})$. By analogy with Sec. 3, if g is a homogeneous polynomial then we denote by T_g the subgraph of T generated by $\text{supp}(g)$.

Let g be a polynomial in $M(X; T)$. Replace all x_n 's in g with λx_{n-1} and represent the obtained polynomial as a linear combination of basis elements. We denote this linear combination by $\bar{\varphi}_{\lambda,p}(g)$. Analogously, denote by $\underline{\varphi}_{\lambda,p}(g)$ the polynomial obtained by replacing all x_n 's in g with λx_{n-2} and representing the obtained polynomial as a linear combination of basis elements. Obviously, $\varphi_{\lambda,p}(g) = \bar{\varphi}_{\lambda,p}(g) + \underline{\varphi}_{\lambda,p}(g) + h$. Here, if $\bar{\varphi}_{\lambda,p}(g) \neq 0$ and $h \neq 0$ then $\text{mdeg}(\bar{\varphi}_{\lambda,p}(g)) > \text{mdeg}(w)$ for any monomial w appearing in h . Similarly, if $\underline{\varphi}_{\lambda,p}(g) \neq 0$ and $h \neq 0$ then $\text{mdeg}(\underline{\varphi}_{\lambda,p}(g)) < \text{mdeg}(w)$ for any monomial w appearing in h . Consequently, if $\varphi_{\lambda,p}(g) = 0$, then $\bar{\varphi}_{\lambda,p}(g) = \underline{\varphi}_{\lambda,p}(g) = 0$.

Lemma 5.6. *For any $g \in M'(X; T) \setminus \{0\}$ there exist positive integers λ_0 and p_0 such that $p_0 = p_0(g)$, $\lambda_0 = \lambda_0(g, p_0)$, and $\varphi_{\lambda,p}(g) \neq 0$ for any $\lambda \geq \lambda_0$ and $p \geq p_0$.*

Proof. By Lemma 5.5, we are left to consider the case when g is homogeneous.

Obviously, if $\varphi_{\lambda,p}(g.y) \neq 0$ for a generator y , then $\varphi_{\lambda,p}(g) \neq 0$.

Let $g \neq 0$. Without loss of generality we may suppose that $\text{supp}(g)$ contains all generators but one of them. Indeed, since T is connected we have $\text{supp}(g) \neq X$ by Theorem 3.3. On the other hand, if $|\text{supp}(g)| < n-1$ then Theorem 4.10 and Theorem 4.12 imply there exists $y \in X \setminus \text{supp}(g)$ such that $g.y \neq 0$. If $|\text{supp}(g.y)| < n-1$, then there exists z , such that $(g.y).z \neq 0$ and so on. Finally, we obtain an element $g' \neq 0$ having all generators but one of them and $\varphi_{\lambda,p}(g') \neq 0$ implies $\varphi_{\lambda,p}(g) \neq 0$.

So, let $|\text{supp}(u)| = n-1$ and $\varphi_{\lambda,p}(g) = 0$. There are several cases possible.

1. If $\text{mdeg}_n(g) = 0$, namely if $x_n \notin \text{supp}(g)$, then we have $\varphi_{\lambda,p}(g) = g \neq 0$ for any λ, p .

2. Let $x \notin \text{supp}(g)$, then $x_{n-1}, x_{n-2} \in \text{supp}(g)$. It is obvious that T_g is a forest. Denote by $T_0, T_1, \dots, T_r$ its connected components, where $T_0 = \{x_n\}$, and $y_1 = x_{n-1}, y_2 = x_{n-2}, y_3, \dots, y_r$ are the greatest vertices of $T_1, T_2, \dots, T_r$ respectively. Note that if v is a monomial appearing in g then the subgraph of T generated by $\text{supp}(\varphi_{\lambda,p}(g))$ is of the form $\bigcup_{i=1}^r T_i$. In particular its connected components are $T_1, T_2, \dots, T_r$.

2.1. Let the smallest vertex of T_g be in T_1 . Since $n \geq 4$, T_g has at least three vertices. Therefore, the smallest vertex of T_g is not equal to x_{n-1} . We have

$$\begin{aligned} g &= \alpha_0 w(x_n, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n) + \alpha_2 w(x_{n-2}, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n) \\ &\quad + \sum_{i=3}^r \alpha_i w(y_i, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n). \end{aligned}$$

Hence

$$(31) \quad \begin{aligned} \underline{\varphi}_{\lambda,p}(g) &= \lambda^{\delta_n}(\alpha_0 + \alpha_2)w'(x_{n-2}, \delta(n-3), \delta_{n-2} + \delta_n, \delta_{n-1}) \\ &\quad + \lambda^{\delta_n} \sum_{i=3}^r \alpha_i w'(y_i, \delta(n-3), \delta_{n-2} + \delta_n, \delta_{n-1}); \end{aligned}$$

$$(32) \quad \begin{aligned} \overline{\varphi}_{\lambda,p}(g) &= \lambda^{\delta_n} p^{\delta_n} \alpha_2 w'(x_{n-2}, \delta(n-3), \delta_{n-2}, \delta_{n-1} + \delta_n) \\ &\quad + \lambda^{\delta_n} p^{\delta_n} \sum_{i=3}^r \alpha_i w'(y_i, \delta(n-3), \delta_{n-2}, \delta_{n-1} + \delta_n). \end{aligned}$$

Thus, the right-hand sides of (31) and (32) are linear combinations of basis elements. Therefore, $\alpha_i = 0$ for $i = 2, 3, \dots, r$ and $\alpha_0 + \alpha_2 = 0$. Consequently, $\alpha_0 = 0$ as well. So, $g = 0$, we get a contradiction.

2.2. Let the smallest vertex of T_g be in T_2 . We have

$$\begin{aligned} g &= \alpha_0 w(x_n, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n) + \alpha_1 w(x_{n-1}, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n) \\ &\quad + \sum_{i=3}^r \alpha_i w(y_i, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n). \end{aligned}$$

Hence,

$$(33) \quad \begin{aligned} \underline{\varphi}_{\lambda,p}(g) &= \lambda^{\delta_n} \alpha_1 w'(x_{n-1}, \delta(n-3), \delta_{n-2} + \delta_n, \delta_{n-1}) \\ &\quad + \lambda^{\delta_n} \sum_{i=3}^r \alpha_i w'(y_i, \delta(n-3), \delta_{n-2} + \delta_n, \delta_{n-1}); \end{aligned}$$

$$(34) \quad \begin{aligned} \overline{\varphi}_{\lambda,p}(g) &= \lambda^{\delta_n} p^{\delta_n} (\alpha_0 + \alpha_1) w'(x_{n-2}, \delta(n-3), \delta_{n-2}, \delta_{n-1} + \delta_n) \\ &\quad + \lambda^{\delta_n} p^{\delta_n} \sum_{i=3}^r \alpha_i w'(y_i, \delta(n-3), \delta_{n-2}, \delta_{n-1} + \delta_n). \end{aligned}$$

The right-hand sides of (33) and (34) are linear combinations of basis elements. Consequently, $\alpha_i = 0$ for $i = 1, 3, 4, \dots, r$ and $\alpha_0 + \alpha_1 = 0$. Therefore, $\alpha_0 = 0$. So, $g = 0$. It is again a contradiction.


2.3. Suppose that the smallest vertex of T_g is neither in T_1 nor in T_2 . Without loss of generality we may suppose that it is in T_r . We have

$$\begin{aligned} g &= \alpha_0 w(x_n, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n) + \alpha_1 w(x_{n-1}, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n) \\ &\quad + \alpha_2 w(x_{n-2}, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n) + \sum_{i=3}^{r-1} \alpha_i w(y_i, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n). \end{aligned}$$

Therefore,

$$(35) \quad \begin{aligned} \underline{\varphi}_{\lambda,p}(g) &= \lambda^{\delta_n} \alpha_1 w'(x_{n-1}, \delta(n-3), \delta_{n-2} + \delta_n, \delta_{n-1}) \\ &\quad + \lambda^{\delta_n} (\alpha_0 + \alpha_2) w'(x_{n-2}, \delta(n-3), \delta_{n-2} + \delta_n, \delta_{n-1}) \\ &\quad + \lambda^{\delta_n} \sum_{i=3}^{r-1} \alpha_i w'(y_i, \delta(n-3), \delta_{n-2} + \delta_n, \delta_{n-1}); \end{aligned}$$

$$(36) \quad \begin{aligned} \overline{\varphi}_{\lambda,p}(g) &= \lambda^{\delta_n} p^{\delta_n} (\alpha_0 + \alpha_1) w'(x_{n-1}, \delta(n-3), \delta_{n-2}, \delta_{n-1} + \delta_n) \\ &\quad + \lambda^{\delta_n} p^{\delta_n} \alpha_2 w'(x_{n-2}, \delta(n-3), \delta_{n-2}, \delta_{n-1} + \delta_n) \\ &\quad + \lambda^{\delta_n} p^{\delta_n} \sum_{i=3}^{r-1} \alpha_i w'(y_i, \delta(n-3), \delta_{n-2}, \delta_{n-1} + \delta_n). \end{aligned}$$

FIGURE 3. Graphs T' and T^* .

The right-hand sides of (35) and (36) are linear combinations of basis elements. Hence, $\alpha_i = 0$ for $i = 1, 2, \dots, r-1$ and $\alpha_0 + \alpha_1 = 0$. Consequently, $\alpha_0 = 0$ as well. As in two last cases, $g = 0$, a contradiction.

3. Suppose that $x, x_n \in \text{supp}(g)$. Let $T_0, T_1, \dots, T_r$ be connected components of T_g and $y_0 = x_n, y_1, \dots, y_r$ the greatest vertices of the corresponding components. In this case, x is the smallest vertex of $\text{supp}(g)$ and $x \in T_0$. Let $\tilde{T}$ be the subgraph of T generated by $\text{supp}(\varphi_{\lambda,p}(g))$. Then the connected components of $\tilde{T}$ are $T'_0, T_1, T_2, \dots, T_r$, where T'_0 is a subgraph of T_0 by deleting x_n (and the incident edge $\{x_n, x\}$). We obtain the following representation:

$$(37) \quad g = \sum_{i=1}^r \alpha_i w(y_i, \delta(n-3), \delta_{n-2}, \delta_{n-1}, \delta_n).$$

Consequently,

$$(38) \quad \overline{\varphi}_{\lambda,p}(g) = \lambda^{\delta_n} p^{\delta_n} \sum_{i=1}^r \alpha_i w'(y_i, \delta(n-3), \delta_{n-2}, \delta_{n-1} + \delta_n).$$

The right-hand side of (38) is a linear combination of basis elements. Therefore, $\alpha_i = 0$ for $i = 1, 2, \dots, r$. Thus, $g = 0$ and we get a contradiction. $\square$

Let T be a tree with the set of vertices X . We say that an endpoint z of T is *unnecessary* if degree of the vertex adjacent to z is at least 3.

Let X^* denote the set of all vertices of T which are not endpoints and let T^* be a subgraph of T generated by X^* .

Now, we describe the procedure constructing a tree T' with no unnecessary vertices for any tree T . If there are unnecessary vertices in T then we delete one of them as well as the edge incident to it. If the obtained graph still has unnecessary vertices we again delete one of them. Let us continue this procedure until the resulting graph has no unnecessary vertices. Denote by T' the obtained tree and by X' the set of its vertices (Figure 3 gives an example).

It is easy to see that for any graph T the obtained graphs with no unnecessary vertices are isomorphic no matter what vertices are deleted.

Lemma 5.7. *Let L_1 and L_2 be two Lie algebras and let $\varphi_{\lambda,p} : L_1 \rightarrow L_2$ be the set of homomorphisms with integer positive parameters λ and p . Suppose that for any $g \in L_1$ there are integers p_0 and $\lambda_0 = \lambda_0(p_0)$ such that $\varphi_{\lambda,p}(g) \neq 0$ whenever $p \geq p_0$ and $\lambda \geq \lambda_0$. Then for any finite set $\Gamma = \{g_1, \dots, g_m\}$ of elements in L_1 there exist integers λ and p such that the restriction of $\varphi_{\lambda,p}$ on Γ (with partial Lie algebra operations) is an inclusion.*

Proof. Let us extend $G = \{g_1, \dots, g_m\}$ adding the elements $g_i - g_j$, $g_i + g_j - g_k$, $[g_i, g_j] - g_k$ for all $i, j, k = 1, 2, \dots, m$. Denote by $\bar{\Gamma}$ the obtained set. It is sufficient to show that there exist λ and p such that the kernel of $\varphi_{\lambda,p} : L_1 \rightarrow L_2$ is disjoint with $\bar{\Gamma}$. Indeed, if it is the case then the images of the elements in Γ are distinct. Moreover, if $g_i \neq g_j + g_k$ or $g_i \neq [g_j, g_k]$ then the images of g_i and $g_j + g_k$ (images of g_i and $[g_j, g_k]$ respectively) are not equal either.

By Lemmas 5.4 and 5.6, for each $g \in \bar{\Gamma}$ one can choose $p_0(g)$ and $\lambda_0(g, p)$ such that $\varphi_{\lambda,p}(g) \neq 0$ for any $p \geq p_0(g)$ and $\lambda \geq \lambda_0(g, p)$. First, let us choose $p_0(g)$ for each $g \in \bar{\Gamma}$. Denote by p_0 the greatest value out of $p_0(g)$. For this p_0 , we find $\lambda_0(g, p_0)$ for each g . Let λ_0 be the greatest number among them. Then, for any $\lambda \geq \lambda_0$, $p \geq p_0$, and $g \in \bar{\Gamma}$ we obtain $\varphi_{\lambda,p}(g) \neq 0$. The proof is complete. $\square$

Lemma 5.8. $T_1^* \simeq T_2^*$ if and only if $T_1' \simeq T_2'$.

Proof. Note that if T is a tree then so is T^* . The graph T' is obtained from the graph T^* as follows: for each endpoint x of the graph T^* , we add a new vertex y and the edge $\{x, y\}$. Indeed, all vertices of T^* are not endpoints in T . It means that their degrees in T' are at least 2. If an edge of T' is not contained in T^* then it cannot connect two vertices of X^* , otherwise, there is a cycle in T' . Moreover, different endpoints of T' are adjacent to different vertices, because each vertex of X^* cannot be adjacent to more than one endpoint in T' . We are left to notice that if x is not an endpoint in T^* then its degree in T^* is at least 2. It means that if T has an endpoint adjacent to x , then there are no these vertices in T' .

Let $T_1^* \simeq T_2^*$. Since T' can be obtained from T^* uniquely, we have $T_1' \simeq T_2'$.

The inverse statement is obvious, i.e. if $T_1' \simeq T_2'$ then $T_1^* \simeq T_2^*$. $\square$

Theorem 5.9. *Let T_1 and T_2 be trees with the sets of vertices $X = \{x_1, x_2, \dots, x_{n_1}\}$ and $Y = \{y_1, y_2, \dots, y_{n_2}\}$ respectively, $n_1, n_2 \geq 2$. The algebras $M(X; T_1)$ and $M(Y; T_2)$ are universally equivalent if and only if the trees T_1^* and T_2^* are isomorphic.*

Proof. Let the algebras $M(X; T_1)$ and $M(Y; T_2)$ be universally equivalent. We prove that T_1^* and T_2^* are isomorphic.

If all vertices of T_1 are endpoints then $n_1 = 2$. Consequently, $M(X; T_1)$ is the free two-generated abelian Lie algebra. So, the formula

$$(39) \quad \exists v_1 v_2 ([v_1, v_2] \neq 0)$$

does not hold in $M(X; T_1)$. Therefore, this formula does not hold in $M(Y; T_2)$ either. But if $n_2 \geq 3$ then T_2 is not a complete graph (because T_2 is a tree). Consequently, $[y_{i_1}, y_{i_2}] \neq 0$ for some vertices $y_{i_1}, y_{i_2} \in Y$. So, formula (39) holds. We get a contradiction. Therefore, $n_2 = 2$ but all trees with two vertices are isomorphic. It means that $T_1 \simeq T_2$, therefore, $T_1^* \simeq T_2^*$ and we are done. Note that, in this case, T_1^* and T_2^* are empty graphs.

Let each graph T_1 and T_2 have exactly one vertex that is not an endpoint. Then $|X^*| = |Y^*| = 1$ and the graphs T_1^* and T_2^* are isomorphic.

Lemma 5.1 implies that if $m \geq 2$ vertices of T_1 are not endpoints then at least two vertices of T_2 are not endpoints.

By Lemma 5.2, the formula $\Phi(T_1)$ constructed by T_1 (see (29)) holds in $M(X; T_1)$. Since $M(X; T_1)$ and $M(Y; T_2)$ are universally equivalent, $\Phi(T_1)$ also holds in $M(Y; T_2)$. Elements z_i can be written as $z_i = d_i + c_i$, where $d_i = \sum_{j=1}^n \alpha_{i,j} y_j$ for some $\alpha_{i,j} \in \mathbb{Z}$ and $c_i \in M'(Y; T_2)$. Since $[u_i, v_i] \in M'(Y; T_2)$, we obtain $[u_i, v_i, z_i] = [u_i, v_i, d_i + c_i] = [u_i, v_i, d_i]$. Therefore, $[u_i, v_i] \in \mathcal{C}(d_i)$. But if the sum $d_i = \sum_{j=1}^n \alpha_{i,j} y_j$ has at least two non-zero summands, then $\mathcal{C}(d_i) = 0$. On the other hand, if $d_i = 0$ (i.e. $z_i \in M'(Y; T_2)$), then $[u_s, v_s, z_i] = 0$ for each $s \neq i$. Therefore, $z_i = \beta_i y_{k_i} + c_i$, where $\beta_i \neq 0$.

Let us prove some conditions of the elements z_i in $M(Y; T_2)$.

1. If $i \neq j$, then $y_{k_i} \neq y_{k_j}$.

Indeed, if the equality $y_{k_i} = y_{k_j}$ holds for some i and j , then we obtain $0 = [u_i, v_i, z_i] = [u_i, v_i, \beta_i y_{k_i} + c_i] = \beta_i [u_i, v_i, y_{k_i}]$. So, $[u_i, v_i, y_{k_i}] = 0$. On the other hand, using the same arguments for $[u_i, v_i, z_j]$ gives $0 \neq [u_i, v_i, y_{k_j}] = [u_i, v_i, y_{k_i}]$. We get a contradiction.

2. The vertices $y_{k_1}, y_{k_2}, \dots, y_{k_m}$ are not endpoints.

Let y_{k_i} be an endpoint. We have $0 = [u_i, v_i, z_i] = [u_i, v_i, \beta_i y_{k_i} + c_i] = \beta_i [u_i, v_i, y_{k_i}]$. Consequently, $[u_i, v_i] \in \mathcal{C}(y_{k_i})$. So, by Theorem, 4.9 $[u_i, v_i] = 0$. It is a contradiction.

3. If x_i and x_j are adjacent in T_1 , then y_{k_i} and y_{k_j} are adjacent in T_2 .

If there is the edge $\{x_i, x_j\}$ in T_1 , then $[z_i, z_j] = 0$. Therefore, we have in $M(Y; T_2)$:

$$\begin{aligned} (40) \quad 0 &= [\beta_i y_{k_i} + c_i, \beta_j y_{k_j} + c_j] \\ &= \beta_i \beta_j [y_{k_i}, y_{k_j}] + \beta_i [y_{k_i}, c_j] + \beta_j [c_i, y_{k_j}] \end{aligned}$$

Since the lengths of all summands in representations of $[y_{k_i}, c_j]$ and $\beta_j [c_i, y_{k_j}]$ as linear combinations of basis elements are greater than 2, $[y_{k_i}, y_{k_j}] = 0$. This means that y_{k_i} and y_{k_j} are adjacent in T_2 .

Consider the map $\varphi : T_1^* \rightarrow T_2^*$ defined as follows: $\varphi(x_i) = y_{k_i}$. By the properties proved above φ is an injective homomorphism. Interchanging T_1 and T_2 and using the same arguments, we obtain there exists an injective homomorphism $\psi : T_2^* \rightarrow T_1^*$. So, in particular $|X^*| = |Y^*|$. Therefore, φ is an isomorphism of the trees T_1^* and T_2^* .

Conversely, suppose that $T_1^* \simeq T_2^*$. Let us show that $M(X; T_1)$ and $M(Y; T_2)$ are universally equivalent.

Suppose that the tree $\tilde{T}$ is obtained from the tree T by deleting the unnecessary vertex x_n and that $\tilde{X} = X \setminus \{x_n\}$. Let us show that the universal theories of $M(X; T)$ and $M(\tilde{X}; \tilde{T})$ coincide. By Theorem 2.11, we are left to prove that any finite submodel of $M(X; T)$ has an isomorphic submodel in $M(\tilde{X}; \tilde{T})$ and vice versa.

In one direction the statement is obviously true because $M(\tilde{X}; \tilde{T})$ is a subalgebra of $M(X; T)$. Therefore, any finite submodel $\{g_1, \dots, g_m\}$ of $M(\tilde{X}; \tilde{T})$ can be considered as a finite submodel of $M(X; T)$. In the other direction, the statement holds by Lemma 5.7.

So, the universal theories of $M(X; T)$ and $M(\tilde{X}; \tilde{T})$ coincide.

Consider the trees T_1 and T_2 . Deleting unnecessary vertices of these graphs one by one, we obtain the graphs T_1' and T_2' . As we just proved, the algebras $M(X'; T_1')$ and $M(X; T_1)$ are universally equivalent. Analogously, the algebras $M(Y'; T_2')$ and $M(Y; T_2)$ are also universally equivalent. Since $T_1^* \simeq T_2^*$ we have $T_1' \simeq T_2'$ as was shown above. Therefore, the universal theories of $M(X'; T_1')$ and $M(Y'; T_2')$ coincide. Consequently, the algebras $M(X; T_1)$ and $M(Y; T_2)$ are universally equivalent. $\square$

Let us note that partially commutative metabelian groups and their theories were studied in [4, 7, 5]. In [7], the criterium of universal equivalence for partially commutative metabelian groups whose defining graphs are trees was obtained. The obtained there necessary and sufficient conditions for universal equivalence of groups are same as the conditions of Theorem 5.9.

REFERENCES

- [1] Bokut L. A., Basis for free polynilpotent Lie algebras, Algebra and Logic, Seminar, **2**, 4 (1963), 13–20 (in Russian).
- [2] Cartier F., Foata D., Problems combinatorics de computation et rearrangements, Lecture Notes in Mathematics, 85, Springer-Verlag, Berlin, 1969.
- [3] Daniyarova .Yu., Kazachkov I. V., Remeslennikov V.N., Algebraic Geometry over Free Metabelian Lie Algebra I: U -Algebras and Universal Classes, J. of Math. Sci., **135**, 5 (2006), 3292–3310.
- [4] Gupta Ch. K., Timoshenko E. I., Partially Commutative Metabelian Groups: Centralizers and Elementary Equivalence, Algebra and Logic, **48**, 3 (2009), 173–192.
- [5] Gupta Ch. K., Timoshenko E. I., Universal Theories for Partially Commutative Metabelian Groups, Algebra and Logic, **50**, 1 (2011), 3–25.
- [6] Kim K. H., Makar-Limanov L., Neggers J., Roush F. W., Graph Algebras, Journal of algebra, **64** (1980), 46–51.
- [7] Timoshenko E. I., Universal Equivalence of Partially Commutative Metabelian Groups, Algebra and Logic, **49**, 2 (2010), 177–196.
- [8] Shmelkin, A. L., Free polynilpotent groups, Izv. AN. USSR, 28 (1964), 91–122 (in Russian).